

CHEMISTRY 9701
THEORY QUESTIONS
AS: CHEMICAL ENERGETICS
www.megalecture.com

3 Bond energies

3(a) Bond energies in diatomic molecules (these are exact values)

Homonuclear

Bond	Energy/kJ mol ⁻¹
H-H	436
D-D	442
N≡N	944
O=O	496
P≡P	485
S=S	425
F-F	158
Cl-Cl	242
Br-Br	193
I-I	151

Heteronuclear

Bond	Energy/kJ mol ⁻¹
H-F	562
H-Cl	431
H-Br	366
H-I	299
C≡O	1077

3(b) Bond energies in polyatomic molecules (these are average values)

Homonuclear

Bond	Energy/kJ mol ⁻¹
C-C	350
C=C	610
C≡C	840
C \equiv C (benzene)	520
N-N	160
N=N	410
O-O	150
Si-Si	222
P-P	200
S-S	264

Heteronuclear

Bond	Energy/kJ mol ⁻¹
C-H	410
C-Cl	340
C-Br	280
C-I	240
C-N	305
C=N	610
C≡N	890
C-O	360
C=O	740
C=O in CO ₂	805
N-H	390
N-Cl	310
O-H	460
Si-Cl	359
Si-H	320
Si-O (in SiO ₂ (s))	460
Si=O (in SiO ₂ (g))	640
P-H	320
P-Cl	330
P-O	340
P=O	540
S-H	347
S-Cl	250
S-O	360
S=O	500

CHEMICAL ENERGETICS (2002-2014)

(b) Carbon disulfide is readily combusted to give CO_2 and SO_2 .

(i) Construct a balanced equation for the complete combustion of CS_2 .

.....

(ii) Define the term *standard enthalpy change of combustion*, ΔH_c^\ominus .

.....

.....

.....

[3]

(c) Calculate the standard enthalpy change of formation of CS_2 from the following data. Include a sign in your answer.

standard enthalpy change of combustion of $\text{CS}_2 = -1110 \text{ kJ mol}^{-1}$

standard enthalpy change of formation of $\text{CO}_2 = -395 \text{ kJ mol}^{-1}$

standard enthalpy change of formation of $\text{SO}_2 = -298 \text{ kJ mol}^{-1}$

[3]

s/13/qp23


(c) The standard enthalpy changes of formation of $\text{NH}_3(\text{g})$ and $\text{H}_2\text{O}(\text{g})$ are as follows.

$$\text{NH}_3(\text{g}), \Delta H_f^\circ = -46.0 \text{ kJ mol}^{-1}$$

$$\text{H}_2\text{O}(\text{g}), \Delta H_f^\circ = -242 \text{ kJ mol}^{-1}$$

Use these data and the value of $\Delta H_{\text{reaction}}^\circ$ given below to calculate the standard enthalpy change of formation of $\text{NO}(\text{g})$.

Include a sign in your answer.



[4]

s/13/qp21

- 3 For some chemical reactions, such as the thermal decomposition of potassium hydrogencarbonate, KHCO_3 , the enthalpy change of reaction cannot be measured directly.

In such cases, the use of Hess' Law enables the enthalpy change of reaction to be calculated from the enthalpy changes of other reactions.

- (a) State Hess' Law.

.....
.....
..... [2]

In order to determine the enthalpy change for the thermal decomposition of potassium hydrogencarbonate, two separate experiments were carried out.

experiment 1

30.0 cm³ of 2.00 mol dm⁻³ hydrochloric acid (an excess) was placed in a conical flask and the temperature recorded as 21.0 °C.

When 0.0200 mol of potassium carbonate, K_2CO_3 , was added to the acid and the mixture stirred with a thermometer, the maximum temperature recorded was 26.2 °C.

- (b) (i) Construct a balanced equation for this reaction.

.....

- (ii) Calculate the quantity of heat produced in **experiment 1**, stating your units. Use relevant data from the *Data Booklet* and assume that all solutions have the same specific heat capacity as water.

- (iii) Use your answer to (ii) to calculate the enthalpy change per mole of K_2CO_3 . Give your answer in kJ mol⁻¹ and include a sign in your answer.

- (iv) Explain why the hydrochloric acid must be in an excess.

.....
..... [4]

experiment 2

The experiment was repeated with 0.0200 mol of potassium hydrogencarbonate, KHCO_3 . All other conditions were the same.

In the second experiment, the temperature fell from 21.0°C to 17.3°C .

(c) (i) Construct a balanced equation for this reaction.


.....

(ii) Calculate the quantity of heat absorbed in **experiment 2**.

(iii) Use your answer to (ii) to calculate the enthalpy change per mole of KHCO_3 . Give your answer in kJ mol^{-1} and include a sign in your answer.

[3]

(d) When KHCO_3 is heated, it decomposes into K_2CO_3 , CO_2 and H_2O .



Use Hess' Law and your answers to (b)(iii) and (c)(iii) to calculate the enthalpy change for this reaction.

Give your answer in kJ mol^{-1} and include a sign in your answer.

[2]

[Total: 11]

w/11/qp21

The unsaturated hydrocarbon, **E**, is obtained by cracking hexane and is important in the chemical industry.

The standard enthalpy change of combustion of **E** is $-2059 \text{ kJ mol}^{-1}$.

(d) Define the term *standard enthalpy change of combustion*.

.....
..... [2]

When 0.47 g of **E** was completely burnt in air, the heat produced raised the temperature of 200 g of water by 27.5°C . Assume no heat losses occurred during this experiment.

(e) (i) Use relevant data from the *Data Booklet* to calculate the amount of heat released in this experiment.

(ii) Use the data above and your answer to (i) to calculate the relative molecular mass, M_r , of **E**.

[4]

(f) Deduce the molecular formula of **E**.

[1]


w/10/qp21

- 3 Alkanes such as methane, CH_4 , undergo few chemical reactions. Methane will, however, react with chlorine but not with iodine.

Relevant standard enthalpy changes of formation for the reaction of methane with chlorine to form chloromethane, CH_3Cl , are given below.


	$\Delta H_f^\ominus/\text{kJ mol}^{-1}$
CH_4	-75
CH_3Cl	-82
HCl	-92

- (a) (i) Use the data to calculate $\Delta H_{\text{reaction}}^\ominus$ for the formation of CH_3Cl .



- (ii) The corresponding reaction with iodine does **not** take place.

Use bond energy data from the *Data Booklet* to calculate a 'theoretical value' for $\Delta H_{\text{reaction}}$ for the following equation.




- (iii) Suggest why this reaction does **not** in fact occur.

.....

[5]


- (c) The energy of activation for the formation of CH_3Cl is 16kJ mol^{-1} .
Use this figure and your answer to (a)(i) to complete the reaction pathway diagram below showing the formation of CH_3Cl from CH_4 and Cl_2 .
Show clearly the intermediate organic species and the final products.
Indicate on your sketch the relevant enthalpy changes and their values.



[4]

w/09/qp21

- 2 Ketene, $\text{C}_2\text{H}_2\text{O}$, is a member of a class of unsaturated organic compounds that is widely used in pharmaceutical research for the synthesis of organic compounds.



ketene

- (c) (i) Define the term *standard enthalpy change of formation*.

.....

.....

.....


- (ii) Use the data below to calculate the standard enthalpy change of formation of ketene.

	$\Delta H^\ominus/\text{kJ mol}^{-1}$
standard enthalpy change of formation of CO_2	-395
standard enthalpy change of combustion of H_2	-286
standard enthalpy change of combustion of $\text{CH}_2=\text{C}=\text{O}$	-1028


[6]

w/08/qp2

(e) Carbon, hydrogen and ethene each burn exothermically in an excess of air.



Use the data to calculate the standard enthalpy change of formation, ΔH_f^\ominus , in kJ mol^{-1} , of ethene at 298 K.



$\Delta H_f^\ominus = \dots\dots\dots \text{ kJ mol}^{-1}$
[3]

w/07/qp2

The unsaturated hydrocarbon **Z** is obtained by cracking hexane and is important in the chemical industry.

The standard enthalpy change of combustion of **Z** is $-2059 \text{ kJ mol}^{-1}$.

(d) Define the term *standard enthalpy change of combustion*.

.....
..... [2]

When 0.47 g of **Z** were completely burnt in air, the heat produced raised the temperature of 200 g of water by 27.5°C .

(e) (i) Calculate the amount of heat released in this experiment.

(ii) Use the data above and your answer to (i) to calculate the relative molecular mass of **Z**.

[4]

(f) Deduce the molecular formula of **Z**.

[1]

w/06/qp2

- (c) Explain the term *standard enthalpy change of formation*, ΔH_f^\ominus .

.....
.....
..... [3]

- (d) Calculate the standard enthalpy change of formation of CS_2 from the following data.

standard enthalpy change of formation of SO_2 = -298 kJ mol^{-1}

standard enthalpy change of formation of CO_2 = -395 kJ mol^{-1}

standard enthalpy change of combustion of CS_2 = $-1110 \text{ kJ mol}^{-1}$


[3]

w/05/qp2

- (d) (i) Explain how enthalpy changes, ΔH values, for covalent bonded molecules can be calculated from bond energies.

.....
.....

- (ii) Use bond energies from the *Data Booklet* to calculate ΔH for the following dissociation.



[3]

Fahad H. Ahmad

- 3 (a) (i) What is meant by the *standard enthalpy change of formation*, ΔH°_f , of a compound? Explain what is meant by the term *standard*.

.....
.....
.....

- (ii) Write an equation, with state symbols, for the ΔH°_f of water.

.....

- (iii) Explain why the ΔH°_f for water is identical to the standard enthalpy change of combustion of hydrogen.

.....

.....[4]

- (b) When calcium is placed in water, aqueous calcium hydroxide is formed and hydrogen is given off.

- (i) Write the equation for the reaction of calcium with water.

.....

- (ii) When 1.00 g of calcium is placed in 200 g of water, the temperature increases by 12.2 °C when the reaction is completed. The specific heat capacity of water, c , is 4.2 J g⁻¹ K⁻¹.

Calculate the heat released in the experiment.

- (iii) Calculate the standard enthalpy change of reaction in kJ mol^{-1} for your equation in (b)(i).

[4]

- (c) (i) State *Hess' Law*.

.....
.....
.....

- (ii) Use Hess' Law and your result in (b)(iii) to calculate the ΔH_f^\ominus of Ca(OH)_2 (aq). You also need the ΔH_f^\ominus of water which is -286 kJ mol^{-1} .

[4]

- (d) Calculate the volume of hydrogen, measured at room temperature and pressure, liberated in the experiment described in (b)(ii).

[2]

[Total : 14]


w/03/qp2

- (d) Most of the ammonia produced which is not used as fertiliser, is oxidised to nitric acid, HNO_3 .

Construct an equation for the oxidation of ammonia by atmospheric oxygen to form nitric acid.

.....[1]

- (e) Urea, $\text{CO}(\text{NH}_2)_2$, is a naturally occurring substance which can be hydrolysed with water to form ammonia according to the following equation.



The standard enthalpy changes of formation of water, urea, carbon dioxide and ammonia (in aqueous solution) are given below.

compound	$\Delta H_f^\ominus / \text{kJ mol}^{-1}$
$\text{H}_2\text{O}(\text{l})$	-287.0
$\text{CO}(\text{NH}_2)_2(\text{aq})$	-320.5
$\text{CO}_2(\text{aq})$	-414.5
$\text{NH}_3(\text{aq})$	-81.0

Use these data to calculate the standard enthalpy change for the hydrolysis of urea.

[2]

w/02/qp2

- 3 With the prospect that fossil fuels will become increasingly scarce in the future, many compounds are being considered for use in internal combustion engines. One of these is DME or dimethyl ether, CH_3OCH_3 . DME is a gas which can be synthesised from methanol. Methanol can be obtained from biomass, such as plant waste from agriculture.

- (a) Define, with the aid of an equation which includes state symbols, the standard enthalpy change of combustion, ΔH_c^\ominus , for DME at 298 K.

equation

definition


.....

..... [3]

- (b) DME may be synthesised from methanol. Relevant enthalpy changes of formation, ΔH_f^\ominus , for this reaction are given in the table below.

compound	$\Delta H_f^\ominus/\text{kJ mol}^{-1}$
$\text{CH}_3\text{OH}(\text{l})$	-239
$\text{CH}_3\text{OCH}_3(\text{g})$	-184
$\text{H}_2\text{O}(\text{l})$	-286

Use these values to calculate $\Delta H_{\text{reaction}}^\ominus$ for the synthesis of DME, using the following equation. Include a sign in your answer.



$$\Delta H_{\text{reaction}}^\ominus = \dots\dots\dots \text{kJ mol}^{-1}$$

[3]

2 Alcohols such as methanol, CH₃OH, are considered to be possible replacements for fossil fuels because they can be used in car engines.

(a) Define, with the aid of an equation which includes state symbols, the standard enthalpy change of combustion, ΔH_c^\ominus , for methanol at 298 K.

equation

definition


.....

..... [3]

Methanol may be synthesised from carbon monoxide and hydrogen. Relevant ΔH_c^\ominus values for this reaction are given in the table below.

compound	$\Delta H_c^\ominus / \text{kJ mol}^{-1}$
CO(g)	-283
H ₂ (g)	-286
CH ₃ OH(g)	-726

(b) Use these values to calculate $\Delta H_{\text{reaction}}^\ominus$ for the synthesis of methanol, using the following equation. Include a sign in your answer.



$$\Delta H_{\text{reaction}}^\ominus = \dots\dots\dots \text{kJ mol}^{-1}$$

[3]

s/12/qp22

- 3 Methanol, CH₃OH, is considered to be a possible alternative to fossil fuels, particularly for use in vehicles.

Methanol can be produced from fossil fuels and from agricultural waste. It can also be synthesised from carbon dioxide and hydrogen.

- (a) Define, with the aid of an equation which includes state symbols, the standard enthalpy change of formation of carbon dioxide.

equation

definition

.....

..... [3]

- (b) Relevant ΔH_f° values for the reaction that synthesises methanol are given in the table.

compound	$\Delta H_f^\circ / \text{kJ mol}^{-1}$
CO ₂ (g)	-394
CH ₃ OH(g)	-201
H ₂ O(g)	-242

- (i) Use these values to calculate $\Delta H_{\text{reaction}}^\circ$ for this synthesis of methanol.

Include a sign in your answer.



$$\Delta H_{\text{reaction}}^\circ = \dots\dots\dots \text{kJ mol}^{-1}$$

- (ii) Suggest one possible environmental advantage of this reaction. Explain your answer.

.....

.....

[5]


s/12/qp21

- 2 Halogenoalkanes have been widely used as aerosol propellants, refrigerants and solvents for many years.

Fluoroethane, $\text{CH}_3\text{CH}_2\text{F}$, has been used as a refrigerant. It may be made by reacting ethene with hydrogen fluoride.

You are to calculate a value for the C–F bond energy in fluoroethane.

- (a) Use relevant bond energies from the *Data Booklet*, and the equation below to calculate a value for the bond energy of the C–F bond.



C–F bond energy = kJ mol^{-1} [4]

- (b) Another halogenoalkane which was used as a refrigerant, and also as an aerosol propellant, is dichlorodifluoromethane, CCl_2F_2 .

State **two** reasons why compounds such as $\text{CH}_3\text{CH}_2\text{F}$ and CCl_2F_2 have been used as aerosol propellants and refrigerants.

.....
..... [2]

s/11/qp22

- 1 Hydrazine, N_2H_4 , can be used as a rocket fuel and is stored as a liquid. It reacts exothermically with oxygen to give only gaseous products.


The enthalpy change of a reaction such as that between hydrazine and oxygen may be calculated by using standard enthalpy changes of formation.

- (a) Define the term *standard enthalpy change of formation*, ΔH_f^\ominus .

.....

 [3]

- (b) Hydrazine reacts with oxygen according to the following equation.



- (i) Use the data in the table to calculate the standard enthalpy change of this reaction.

compound	$\Delta H_f^\ominus/\text{kJ mol}^{-1}$
$\text{N}_2\text{H}_4(\text{l})$	50.6
$\text{H}_2\text{O}(\text{g})$	-241.8

$\Delta H^\ominus = \dots\dots\dots \text{kJ mol}^{-1}$

- (ii) Although the above reaction is highly exothermic, hydrazine does not burn spontaneously in oxygen. Suggest a reason for this.

.....

s/10/qp23


- (ii) Use the *Data Booklet* to calculate the enthalpy change that occurs when one mole of gaseous magnesium ions, Mg^{2+} , is formed from one mole of gaseous magnesium atoms.

Include a sign in your answer.

enthalpy change = kJ mol^{-1}
[3]

s/09/qp2

- (d) The equation for the complete combustion of ethyne is given below.
Use appropriate bond energy data from the *Data Booklet* to calculate a value for the enthalpy change of combustion of ethyne.



[3]

- (e) The value for the standard enthalpy change of combustion of ethyne is $-1300 \text{ kJ mol}^{-1}$.

- (i) Define the term *standard enthalpy change of combustion*.

.....
.....
.....

- (ii) Explain why your answer to (d) does not have the same value as the standard enthalpy change of combustion.


.....
.....

[3]

s/06/qp2

- 2 Ethanol, C₂H₅OH, is a most important industrial chemical and is used as a solvent, a fuel and an intermediate in large scale organic synthesis.


Ethanol is prepared industrially by the reaction of ethene and steam in the presence of a catalyst.



The standard enthalpy change of the reaction can be determined by using the standard enthalpy changes of combustion, ΔH_c^\ominus , at 298 K.

	$\Delta H_c^\ominus / \text{kJ mol}^{-1}$
C ₂ H ₄ (g)	-1411
C ₂ H ₅ OH(l)	-1367

- (a) Calculate the standard enthalpy change for the following reaction.



[2]

- (b) (i) Define the term *standard enthalpy change of combustion*.

.....

- (ii) Explain why the state symbols for water and ethanol given in the equation in (a) have been changed from those quoted in the industrial process.

.....

- (iii) Write the equation for the complete combustion of ethanol.

.....
 [4]

Fahad H. Ahmad