

Q1.

- 7 (i) State $\frac{dx}{d\theta} = 2 - 2\cos 2\theta$ or $\frac{dy}{d\theta} = 2\sin 2\theta$ B1
- Use $\frac{dy}{dx} = \frac{dy}{d\theta} \div \frac{dx}{d\theta}$ M1
- Obtain answer $\frac{dy}{dx} = \frac{2\sin 2\theta}{2-2\cos 2\theta}$ or equivalent A1
- Make relevant use of $\sin 2A$ and $\cos 2A$ formulae (indep.) M1
- Obtain given answer correctly A1
- [5]

- (ii) Substitute $\theta = \frac{1}{4}\pi$ in $\frac{dy}{dx}$ and both parametric equations M1
- Obtain $\frac{dy}{dx} = 1, x = \frac{1}{2}\pi - 1, y = 2$ A1
- Obtain equation $y = x + 1.43$, or any exact equivalent A1√
- [3]

- (iii) State or imply that tangent is horizontal when $\theta = \frac{1}{2}\pi$ or $\frac{3}{2}\pi$ B1
- Obtain a correct pair of x , y or x - or y -coordinates B1
- State correct answers $(\pi, 3)$ and $(3\pi, 3)$ B1
- [3]

Q2.

- 6 (i) State that $\frac{dx}{dt} = 2 + \frac{1}{t}$ or $\frac{dy}{dt} = 1 - \frac{4}{t^2}$, or equivalent B1
- Use $\frac{dy}{dx} = \frac{dy}{dt} \div \frac{dx}{dt}$ M1
- Obtain the given answer A1 3
- (ii) Substitute $t = 1$ in $\frac{dy}{dx}$ and both parametric equations M1
- Obtain $\frac{dy}{dx} = -1$ and coordinates $(2, 5)$ A1
- State equation of tangent in any correct horizontal form e.g. $x + y = 7$ A1√ 3
- (iii) Equate $\frac{dy}{dx}$ to zero and solve for t M1
- Obtain answer $t = 2$ A1
- Obtain answer $y = 4$ A1
- Show by any method (but not via $\frac{d}{dt}(y')$) that this is a minimum point A1 4

Q3.

- 5 (i) Differentiate using chain or quotient rule
 Obtain derivative in any correct form
 Obtain given answer correctly
- (ii) State $\frac{dx}{d\theta} = \sec^2 \theta$, or equivalent
 Use $\frac{dy}{dx} = \frac{dy}{d\theta} \div \frac{dx}{d\theta}$
 Obtain given answer correctly

M1
 A1
 A1 3
 B1
 M1
 A1 3

Page 2	Mark Scheme	Syllabus	Paper
	GCE AS LEVEL – JUNE 2005	9709	2

- (iii) State that $\theta = \frac{\pi}{6}$
 Obtain x-coordinate $1 + \frac{1}{\sqrt{3}}$, or equivalent
 Obtain y-coordinate $\frac{2}{\sqrt{3}}$, or equivalent

B1
 B1
 B1 3

Q4.

- 3 State correct derivative $1 - 2\sin x$
 Equate derivative to zero and solve for x
 Obtain answer $x = \frac{1}{6}\pi$
 Carry out an appropriate method for determining the nature of a stationary point
 Show that $x = \frac{1}{6}\pi$ is a maximum with no errors seen
 Obtain second answer $x = \frac{5}{6}\pi$ in range
 Show this is a minimum point
 (f.t. is on the incorrect derivative $1 + 2\sin x$)

B1
 M1
 A1
 M1
 A1
 A1/
 A1/ 7

Q5.

5	(i) State $2y \frac{dy}{dx}$ as the derivative of y^2	B1	
	State $2y + 2x \frac{dy}{dx}$ or equivalent, as derivative of $2xy$	B1	
	Equate attempted derivative of LHS to zero and set $\frac{dy}{dx}$ equal to zero	M1	
	Obtain given relation $y = -3x$ correctly	A1	4
	[The M1 is dependent on at least one B1 being earned earlier.]		
	(ii) Carry out complete method for finding x^2 or y^2	M1	
	Obtain $x^2 = 1$ or $y^2 = 9$	A1	
	Obtain point $(1, -3)$	A1	
	Obtain second point $(-1, 3)$	A1	4

Q6.

3	(i)	State $\frac{dx}{dt} = 3 + \frac{1}{t-1}$ or $\frac{dy}{dt} = 2t$	B1	
		Use $\frac{dy}{dx} = \frac{dy}{dt} \div \frac{dx}{dt}$	M1	
		Obtain $\frac{dy}{dx}$ in any correct form, e.g. $\frac{2t(t-1)}{3t-2}$	A1	[3]
	(ii)	Equate derivative to 1 and solve for t	M1	
		Obtain roots 2 and $\frac{1}{2}$	A1	
		State or imply that only $t=2$ is admissible c.w.o.	A1	
		Obtain coordinates $(6, 5)$	A1	[4]

Q7.

6	(i)	Use product rule	M1*	
		Obtain correct derivative in any fom, e.g. $(x-1)e^x$	A1	
		Equate derivative to zero and solve for x	M1* (dep)	
		Obtain $x = 1$	A1	
		Obtain $y = -e$	A1	[5]
	(ii)	Carry out a method for determining the nature of a stationary point	M1	
		Show that the point is a minimum point, with no errors seen	A1	[2]

Q8.

- 7 (i) State $2y \frac{dy}{dx}$ as derivative of y^2 , or equivalent B1
State $4y + 4x \frac{dy}{dx}$ as derivative of $4xy$, or equivalent B1
Equate derivative of LHS to zero and solve for $\frac{dy}{dx}$ M1
Obtain given answer correctly A1 [4]
[The M1 is dependent on at least one of the B marks being obtained.]
- (ii) State or imply that the coordinates satisfy $2y - x = 0$ B1
Obtain an equation in x^2 (or y^2) M1
Solve and obtain $x^2 = 4$ (or $y^2 = 1$) A1
State answer (2, 1) A1
State answer (-2, -1) A1 [5]

Q9.

- 4 State $\frac{dx}{d\theta} = 4 \cos \theta$ B1
State $\frac{dy}{d\theta} = 4 \sin 2\theta$, or equivalent B1
Use $\frac{dy}{dx} = \frac{dy}{d\theta} \div \frac{dx}{d\theta}$ M1
Obtain $\frac{dy}{dx}$ in any correct form, e.g. $\frac{\sin 2\theta}{\cos \theta}$ A1
Simplify and obtain answer $2 \sin \theta$ A1√ [5]
[The f.t. is on gradients of the form $k \sin 2\theta / \cos \theta$, or equivalent.]

Q10.

- 6 (i) State $2xy + x^2 \frac{dy}{dx}$ as derivative of x^2y B1
State $2y \frac{dy}{dx}$ as derivative of y^2 B1
Equate derivatives of LHS and RHS, and solve for $\frac{dy}{dx}$ M1
Obtain given answer A1 [4]
- (ii) Substitute and obtain gradient $\frac{2}{5}$, or equivalent B1
Form equation of tangent at the given point (1, 2) M1
Obtain answer $2x - 5y + 8 = 0$, or equivalent A1 [3]
[The M1 is dependent on at least one of the B marks being obtained.]

Q11.

- 5 (i) Use product rule M1
Obtain correct derivative in any form A1
Show that derivative is equal to zero when $x = 3$ A1 [3]
- (ii) Substitute $x = 1$ into gradient function, obtaining $2e^{-1}$ or equivalent M1
State or imply required y -coordinate is e^{-1} B1
Form equation of line through $(1, e^{-1})$ with gradient found (NOT the normal) M1
Obtain equation in any correct form A1 [4]

Q12.

- 2 State $\frac{dx}{dt} = 3 + 2 \cos 2t$ or $\frac{dy}{dt} = -4 \sin 2t$ (or both) B1
- Use $\frac{dy}{dx} = \frac{dy}{dt} \div \frac{dx}{dt}$ M1
- Obtain or imply $\frac{-4 \sin 2t}{3 + 2 \cos 2t}$ A1
- Substitute $\frac{1}{6}\pi$ to obtain $-\frac{1}{2}\sqrt{3}$ or exact equivalent A1 [4]

Q13.

- 5 (i) Differentiate $\ln(x-3)$ to obtain $\frac{1}{x-3}$ B1
Attempt to use product rule M1
Obtain $\ln(x-3) + \frac{x}{x-3}$ or equivalent A1
Substitute 4 to obtain 4 A1 [4]
- (ii) Use correct quotient or product rule M1
Obtain correct derivative in any form, e.g. $\frac{(x+1)-(x-1)}{(x+1)^2}$ A1
Substitute 4 to obtain $\frac{2}{25}$ A1 [3]

Q14.

- 5 Obtain $4y \frac{dy}{dx}$ as derivative of $2y^2$ B1
- Differentiate LHS term by term to obtain expression including at least one $\frac{dy}{dx}$ M1
- Obtain $2x + 4y \frac{dy}{dx} + 5 + 6 \frac{dy}{dx}$ A1
- Substitute 2 and -1 to attempt value of $\frac{dy}{dx}$ M1
- Obtain $-\frac{9}{2}$ A1
- Obtain equation $9x + 2y - 16 = 0$ or equivalent of required form A1 [6]

Q15.

- 6 (i) Attempt differentiation using product rule M1
 Obtain $8x \ln x + 4x$ (a.c.f.) A1
 Equate first derivative to zero and attempt solution M1
 Obtain 0.607 A1
 Obtain -0.736 following their x -coordinate A1✓ [5]
- (ii) Use an appropriate method for determining nature of stationary point M1
 Conclude point is a minimum (with no errors seen, second derivative = 8) A1 [2]

Q16.

- 5 (i) State $\frac{dx}{dt} = \frac{1}{t+1}$ B1
 State $\frac{dy}{dt} = 2e^{2t} + 2$ B1
 Attempt expression for $\frac{dy}{dx}$ M1
 Obtain $\frac{dy}{dx} = (2e^{2t} + 2)(t + 1)$ or equivalent A1 [4]
- (ii) Substitute $t = 0$ and attempt gradient of normal M1
 Obtain $-\frac{1}{4}$ following their expression for $\frac{dy}{dx}$ A1✓
 Attempt to find equation of normal through point $(0, 1)$ M1
 Obtain $x + 4y - 4 = 0$ A1 [4]

Q17.

- 5 (i) Use product rule to differentiate y M1
 Obtain correct derivative in any form A1
 Use $\frac{dy}{dx} = \frac{dy}{dt} \div \frac{dx}{dt}$ M1
 Obtain given answer correctly A1 [4]
- (ii) Substitute $t = 0$ in $\frac{dy}{dx}$ and both parametric equations B1
 Obtain $\frac{dy}{dx} = 2$ and coordinates $(1, 0)$ B1
 Form equation of the normal at their point, using negative reciprocal of their $\frac{dy}{dx}$ M1
 State correct equation of normal $y = -\frac{1}{2}x + \frac{1}{2}$ or equivalent A1 [4]

Q18.

- 5 (i) State $3 \frac{dy}{dx}$ as derivative of $3y$, or equivalent B1
- State $4xy + 2x^2 \frac{dy}{dx}$ as a derivative of $2x^2y$, or equivalent B1
- Equate derivative of LHS to zero and solve for $\frac{dy}{dx}$ M1
- Obtain given answer correctly A1 [4]
- (ii) Substitute $x = 2$ into given equation and solve for y M1
- Obtain gradient = $\frac{12}{5}$ correctly A1
- Form equation of the normal at their point, using negative recip of their $\frac{dy}{dx}$ M1
- State correct equation of normal $5x + 12y + 2 = 0$ or equivalent A1 [4]

Q19.

- 7 (i) State $6y \frac{dy}{dx}$ as the derivative of $3y^2$ B1
- State $\pm 2x \frac{dy}{dx} \pm 2y$ as the derivative of $-2xy$ (allow any combination of signs here) B1
- Equate attempted derivative of LHS to 0 (or 10) and solve for $\frac{dy}{dx}$ M1
- Obtain the given answer correctly A1 4
- [The M1 is dependent on at least one of the B marks being earned.]
- (ii) State or imply the points lie on $y - 2x = 0$ or $(y - 2x) / (3y - 2x) = 0$ B1 ⊕
- Carry out complete method for finding one coordinate of a point of intersection of $y = kx$ with the given curve M1
- Obtain $10x^2 = 10$ or $2\frac{1}{2}y^2 = 10$ or 2-term equivalent A1
- Obtain one correct point e.g. (1, 2) or 2 values of x (or y) A1
- Obtain a second correct point e.g. (-1, -2) A1 $\sqrt{}$ 5 ⊕

Q20.

6 (i)	State A is $(4, 0)$ State B is $(0, 4)$	B1 B1	
			[2]
(ii)	Use the product rule to obtain the first derivative Obtain derivative $(4 - x)e^x - e^x$, or equivalent Equate derivative to zero and solve for x Obtain answer $x = 3$ only	M1(dep) A1 M1 (dep) A1	
			[4]
(iii)	Attempt to form an equation in p e.g. by equating gradients of OP and the tangent at P , or by substituting $(0, 0)$ in the equation of the tangent at P Obtain equation in any correct form e.g. $\frac{4-p}{p} = 3 - p$ Obtain 3-term quadratic $p^2 - 4p + 4 = 0$, or equivalent Attempt to solve a quadratic equation in p Obtain answer $p = 2$ only	M1 A1 A1 M1 A1	
			[5]

Q21.

5 (i)	Use the product rule to obtain the first derivative (must involve 2 terms) Obtain derivative $2x \ln x + x^2 \frac{1}{x}$ or equivalent Equate derivative to zero and solve for x Obtain answer $x = e^{-0.5}$ or $\frac{1}{\sqrt{e}}$ or equivalent (e.g. 0.61)	M1 A1 M1 A1	4
(ii)	Determine nature of stationary point using correct second derivative $(3 + 2\ln x)$ or correct first derivative or equation of the curve $(3 y\text{-values, central one } y(\exp(-0.5)))$ Show point is a minimum completely correctly	M1 A1	2

Q22.

4 (i)	State $3y^3 \frac{dy}{dx}$ as derivative of y^3 State $9y + 9x \frac{dy}{dx}$ as derivative of $9xy$ Express $\frac{dy}{dx}$ in terms of x and y Obtain given answer correctly [The M1 is conditional on at least one B mark being obtained.]	B1 B1 M1 A1	4
(ii)	Obtain gradient at $(2, 4)$ in any correct unsimplified form Form the equation of the tangent at $(2, 4)$ Obtain answer $5y - 4x = 12$, or equivalent	B1 M1 A1	3

Q23.

- | | | | |
|---|---|-----|-----|
| 4 | State derivative $2 - \sec^2 x$, or equivalent | B1 | |
| | Equate derivative to zero and solve for x | M1 | |
| | Obtain $x = \frac{1}{4}\pi$, or 0.785 ($\pm 45^\circ$ gains A1) | A1 | |
| | Obtain $x = -\frac{1}{4}\pi$, (allow negative of first solution) | A1√ | |
| | Obtain corresponding y -values $\frac{1}{2}\pi - 1$ and $-\frac{1}{2}\pi + 1$, ± 0.571 | A1 | [5] |

Q24.

- | | | | |
|---|---|-----|-----|
| 6 | At any stage, state the correct derivative of $e^{-\frac{1}{2}x}$ or $e^{\frac{1}{2}x}$ | B1 | |
| | Use product or quotient rule | M1 | |
| | Obtain correct first derivative in any form | A1 | |
| | Obtain correct second derivative in any form | B1√ | |
| | Equate second derivative to zero and solve for x | M1 | |
| | Obtain $x = 4$ | A1 | |
| | Obtain $y = 4e^{-2}$, or equivalent | A1 | [7] |

Q25.

- | | | | | |
|---|------|--|----------|-----|
| 6 | (i) | Use product rule | M1* | |
| | | Obtain derivative in any correct form | A1 | |
| | | Equate derivative to zero and solve for x | M1(dep*) | |
| | | Obtain $x = 1/e$, or exact equivalent | A1 | |
| | | Obtain $y = -1/e$, or exact equivalent | A1 | [5] |
| | (ii) | Carry out complete method for determining the nature of a stationary point | M1 | |
| | | Show that at $x = 1/e$ there is a minimum point, with no errors seen | A1 | [2] |

Q26.

- | | | | | |
|---|------|---|---------|-----|
| 8 | (i) | EITHER: Substitute $x = 1$ and attempt to solve 3-term quadratic in y | M1 | |
| | | Obtain answers $(1, 1)$ and $(1, -3)$ | A1 | |
| | | OR: State answers $(1, 1)$ and $(1, -3)$ | B1 + B1 | [2] |
| | (ii) | State $2y \frac{dy}{dx}$ as derivative of y^2 | B1 | |
| | | State $2y + 2x \frac{dy}{dx}$ as derivative of $2xy$ | B1 | |
| | | Substitute for x and y , and solve for $\frac{dy}{dx}$ | M1 | |
| | | Obtain $\frac{dy}{dx} = 0$ when $x = 1$ and $y = 1$ | A1 | |
| | | Obtain $\frac{dy}{dx} = -2$ when $x = 1$ and $y = -3$ | A1√ | |
| | | Form the equation of the tangent at $(1, -3)$ | M1 | |
| | | Obtain answer $2x + y + 1 = 0$ | A1 | [7] |

Q27.

- 4 (i) State $\frac{dx}{dt} = e^{-t}$ or $\frac{dy}{dt} = e^t - e^{-t}$ B1
 Use $\frac{dy}{dx} = \frac{dy}{dt} \div \frac{dx}{dt}$ M1
 Obtain given answer correctly A1 [3]
- (ii) Substitute $\frac{dy}{dx} = 2$ and use correct method for solving an equation of the form $e^{2t} = a$,
 where $a > 0$ M1
 Obtain answer $t = \frac{1}{2} \ln 3$, or equivalent A1 [2]

Q28.

- 4 (i) State $\frac{dx}{dt} = \frac{1}{t-2}$ or $\frac{dy}{dt} = 1 - 9t^{-2}$ B1
 Use $\frac{dy}{dx} = \frac{dy}{dt} \div \frac{dx}{dt}$ M1
 Obtain given answer correctly A1 [3]
- (ii) Equate derivative to zero and solve for t M1
 State or imply that $t = 3$ is admissible c.w.o., and note $t = -3, 2$ cases A1
 Obtain coordinates (1, 6) and no others A1 [3]

Q29.

- 8 (i) State $2y \frac{dy}{dx}$ as derivative of y^2 , or equivalent B1
 State $2y + 2x \frac{dy}{dx}$ as derivative of $2xy$, or equivalent B1
 Substitute $x = -2$ and $y = 2$ and evaluate $\frac{dy}{dx}$ M1
 Obtain zero correctly and make correct conclusion A1 [4]
- (ii) Substitute $x = -2$ into given equation and solve M1
 Obtain $y = -6$ correctly A1
 Obtain $\frac{dy}{dx} = 2$ correctly B1
 Form the equation of the tangent at $(-2, -6)$ M1
 Obtain answer $y = 2x - 2$ A1 [5]

Q30.

- 3 Obtain derivative of the form $k \sec^2 2x$, where $k = 1$ or $k = \frac{1}{2}$ M1
 Obtain correct derivative $\sec^2 2x$ A1
 Use correct method for solving $\sec^2 2x = 4$ M1
 Obtain answer $x = \frac{1}{6}\pi$ (or 0.524 radians) A1
 Obtain answer $x = \frac{1}{3}\pi$ (or 1.05 radians) and no others in range A1 [5]

Q31.

- 7 (i) Use product rule to differentiate y M1
 Obtain correct derivative in any form in t for y A1
 Use $\frac{dy}{dx} = \frac{dy}{dt} \div \frac{dx}{dt}$ M1
 Obtain given answer correctly A1 [4]
- (ii) State $t = 0$ M1
 State that $\frac{dy}{dx} = 0$ and make correct conclusion A1 [2]
- (iii) Substitute $t = -2$ into equation for x or y M1
 Obtain $(e^{-6}, 4e^{-2} + 3)$ A1 [2]

Q32.

- 6 (i) State $\frac{dx}{dt} = 4 \sin \theta \cos \theta$ or equivalent (nothing for $\frac{dy}{dx} = 4 \sec^2 \theta$) B1
 Use $\frac{dy}{dx} = \frac{dy}{d\theta} \div \frac{dx}{d\theta}$ M1
 Obtain given answer correctly A1 [3]
- (ii) Substitute $\theta = \frac{\pi}{4}$ in $\frac{dy}{dx}$ and both parametric equations M1
 Obtain $\frac{dy}{dx} = 4$ and coordinates $(2, 4)$ A1
 Form equation of tangent at their point M1
 State equation of tangent in correct form $y = 4x - 4$ A1 [4]

Q33.

- 1 Obtain derivative of the form $\frac{k}{5x+1}$, where $k = 1, 5$ or $\frac{1}{5}$ M1
 Obtain correct derivative $\frac{5}{5x+1}$ A1
 Substitute $x = 4$ into expression for derivative and obtain $\frac{5}{21}$ A1√ [3]

Q34.

- 8 (i) State $2y \frac{dy}{dx}$ as derivative of y^2 , or equivalent B1
- Equate derivative of LHS to zero and solve for $\frac{dy}{dx}$ M1
- Obtain given answer correctly A1 [3]
- (ii) Equate gradient expression to -1 and rearrange M1
- Obtain $y = 2x$ A1
- Substitute into original equation to obtain an equation in x^2 (or y^2) M1
- Obtain $2x^2 - 3x - 2 = 0$ (or $y^2 - 3y - 4 = 0$) A1
- Correct method to solve their quadratic equation M1
- State answers $(-\frac{1}{2}, -1)$ and $(2, 4)$ A1 [6]

Q35.

- 4 (i) State $\frac{dx}{dt} = \frac{-2}{1-2t}$ or $\frac{dy}{dt} = -2t^{-2}$ B1
- Use $\frac{dy}{dx} = \frac{dy}{dt} \div \frac{dx}{dt}$ M1
- Obtain given answer correctly A1 [3]
- (ii) Equate derivative to 3 and solve for t M1
- State or imply that $t = -1$ c.w.o. A1
- Obtain coordinates $(\ln 3, -2)$ A1 [3]

Q36.

- 2 Use quotient rule or product rule, correctly M1
- Obtain correct derivative in any form A1
- Equate derivative to zero and solve for x M1
- Obtain $x = \frac{\pi}{8}$ A1 [4]

Q37.

- 7 (i) State $4y \frac{dy}{dx}$ as derivative of $2y^2$, or equivalent B1
 State $4y + 4x \frac{dy}{dx}$ as derivative of $4xy$, or equivalent B1
 Equate derivative of LHS to zero and solve for $\frac{dy}{dx}$ M1
 Obtain given answer correctly A1 [4]
- (ii) State or imply that the coordinates satisfy $3x - 2y = 0$ B1
 Obtain an equation in x^2 (or y^2) M1
 Solve and obtain $x^2 = 4$ (or $y^2 = 9$) A1
 State answer (2, 3) A1
 State answer (-2, -3) A1 [5]

Q38.

- 3 Obtain derivative $e^{2x} - 5e^x + 4$ B1
 Equate derivative to zero and carry out recognisable solution method for a quadratic in e^x M1
 Obtain $e^x = 1$ or $e^x = 4$ A1
 Obtain $x = 0$ and $x = \ln 4$ A1
 Use an appropriate method for determining nature of at least one stationary point M1
 $\left(\frac{d^2y}{dx^2} = 2e^{2x} - 5e^x, \text{ when } x = 0, \frac{d^2y}{dx^2} = -(3), x = \ln 4, \frac{d^2y}{dx^2} = +(12) \right)$
 Conclude maximum at $x = 0$ and minimum at $x = \ln 4$ (no errors seen) A1 [6]

Q39.

- 5 (i) State $\frac{dx}{d\theta} = -2 \sin 2\theta + \sin \theta$ or $\frac{dy}{d\theta} = 8 \sin \theta \cos \theta$ B1
 Use $\frac{dy}{dx} = \frac{dy}{d\theta} \div \frac{dx}{d\theta}$ M1
 Use $\sin 2\theta = 2 \sin \theta \cos \theta$ M1
 Obtain given answer correctly A1 [4]
- (ii) Equate derivative to -4 and solve for $\cos \theta$ M1
 Obtain $\cos \theta = \frac{1}{2}$ A1
 Obtain $x = -1$ A1
 Obtain $y = 3$ A1 [4]

Q40.

- 2 Use quotient or product rule M1
 Obtain correct derivative in any form A1
 Equate (numerator) of derivative to zero and solve for x DM1
 Obtain $x = \frac{1}{3}$ A1
 Obtain $y = \frac{3}{2}$ A1 [5]

Q41.

- 5 (i) State $\frac{dx}{dt} = \frac{1}{2}t^{-\frac{1}{2}}$ or $\frac{dy}{dt} = \frac{3}{t}$ B1
 Use $\frac{dy}{dx} = \frac{dy}{dt} \div \frac{dx}{dt}$ M1
 Use $y = 6$ to find t M1
 Obtain $t = e^2$ A1
 Obtain $\frac{dy}{dx} = \frac{6}{e}$ A1 [5]
- (ii) Obtain x and form equation of the tangent at their point M1
 Obtain correct equation for tangent $\left(y - 6 = \frac{6}{e}(x - (1 + e))\right)$ A1
 Show that tangent passes through $(1, 0)$ by substitution A1 [3]

Q42.

- 2 (i) Differentiate to obtain form $k_1 \cos x + k_2 \sec^2 2x$ M1
 Obtain correct second term $2 \sec^2 2x$ A1
 Obtain $3 \cos x + 2 \sec^2 2x$ and hence answer 5 A1 [3]
- (ii) Differentiate to obtain form $k e^{2x} (1 + e^{2x})^{-2}$ M1
 Obtain correct $-12e^{2x} (1 + e^{2x})^{-2}$ or equivalent (may be implied) A1
 Obtain -3 A1 [3]

Q43.

- 7 (i) Obtain $3y + 3x \frac{dy}{dx}$ as derivative of $3xy$ B1
 Obtain $2y \frac{dy}{dx}$ as derivative of y^2 B1
 State $4x + 3y + 3x \frac{dy}{dx} + 2y \frac{dy}{dx} = 0$ B1
 Substitute 2 and -1 to find gradient of curve (dependent on at least one B1) M1
 Form equation of tangent through $(2, -1)$ with numerical gradient
 (dependent on previous M1) DM1
 Obtain $5x + 4y - 6 = 0$ or equivalent of required form A1 [6]
- (ii) Use $\frac{dy}{dx} = 0$ to find relation between x and y M1
 (dependent on at least one B1 from part(i)) A1
 Obtain $4x + 3y = 0$ or equivalent M1
 Substitute for x or y in equation of curve A1 [4]
 Obtain $-\frac{1}{8}y^2 = 3$ or $-\frac{2}{9}x^2 = 3$ or equivalent and conclude appropriately

Q44.

- 4 Obtain $\frac{dx}{dt} = \frac{2}{t+1}$ B1
 Obtain $\frac{dy}{dt} = 4e^t$ B1
 Use $\frac{dy}{dx} = \frac{dy}{dt} / \frac{dx}{dt}$ with $t = 0$ to find gradient M1
 Obtain 2 A1
 Form equation of tangent through (0, 4) with numerical gradient obtained from attempt to differentiate M1
 Obtain $2x - y + 4 = 0$ or equivalent of required form A1 [6]

Q45.

- 8 (i) Differentiate using product rule M1
 Obtain $\sec^2 x \cos 2x - 2 \tan x \sin 2x$ A1
 Use $\cos 2x = 2 \cos^2 x - 1$ or $\sin 2x = 2 \sin x \cos x$ or both B1
 Express derivative in terms of $\sec x$ and $\cos x$ only M1
 Obtain $4 \cos^2 x - \sec^2 x - 2$ with no errors seen (AG) A1 [5]
- (ii) State $4 \cos^4 x - 2 \cos^2 x - 1 = 0$ B1
 Apply quadratic formula to a 3 term quadratic equation in terms of $\cos^2 x$ to find the least positive value of $\cos^2 x$ M1
 Obtain or imply $\cos^2 x = \frac{1 + \sqrt{5}}{4}$ or 0.809... A1
 Obtain 0.45 A1 [4]

Q46.

- 4 (i) Differentiate to obtain form $k_1 \sin 2x + k_2 \cos x$ M1
 Obtain correct $-6 \sin 2x - 5 \cos x$ A1
 Substitute $\frac{1}{6} \pi$ to obtain $-\frac{11}{2} \sqrt{3}$ or exact equivalent A1 [3]
- (ii) Obtain $6y + 6x \frac{dy}{dx}$ as derivative of $6xy$ B1
 Obtain $3y^2 \frac{dy}{dx}$ as derivative of y^3 B1
 Obtain $3x^2 + 6y + 6x \frac{dy}{dx} + 3y^2 \frac{dy}{dx} = 0$ or equivalent B1
 Substitute 1 and 2 to find value of gradient dependent on at least one B1 M1
 Obtain gradient $-\frac{15}{18}$ or $-\frac{5}{6}$ A1 [5]

Q47.

- | | | |
|---|---|--------|
| 3 | Obtain $6y + 6x \frac{dy}{dx}$ as derivative of $6xy$ | B1 |
| | Obtain $2y \frac{dy}{dx}$ as derivative of y^2 | B1 |
| | Obtain $\frac{3}{x}$ and $\frac{d}{dx}(16) = 0$ | B1 |
| | Substitute 1 and 2 to find value of $\frac{dy}{dx}$ | M1 |
| | Obtain value $\frac{2}{3}$ as gradient of normal following their value of $\frac{dy}{dx}$ | A1✓ |
| | Form equation of normal through (1, 2) with numerical gradient | M1 |
| | Obtain $2x - 3y + 4 = 0$ | A1 [7] |

P3 (variant1 and 3)

Q1.

- | | | |
|---|---|---|
| 9 | <p>(i) Use quotient or product rule to differentiate $(1 - x)/(1 + x)$</p> <p>Obtain correct derivative in any form</p> <p>Use chain rule to find $\frac{dy}{dx}$</p> <p>Obtain a correct expression in any form</p> <p>Obtain the gradient of the normal in the given form correctly</p> | <p>M1</p> <p>A1</p> <p>M1</p> <p>A1</p> <p>A1 [5]</p> |
| | <p>(ii) Use product rule</p> <p>Obtain correct derivative in any form</p> <p>Equate derivative to zero and solve for x</p> <p>Obtain $x = \frac{1}{2}$</p> | <p>M1</p> <p>A1</p> <p>M1</p> <p>A1 [4]</p> |

Q2.

- | | | |
|---|--|-------------------------|
| 2 | <p>(i) Obtain $\frac{k \cos 2x}{1 + \sin 2x}$ for any non-zero constant k</p> <p>Obtain $\frac{2 \cos 2x}{1 + \sin 2x}$</p> | <p>M1</p> <p>A1 [2]</p> |
| | <p>(ii) Use correct quotient or product rule</p> <p>Obtain $\frac{x \sec^2 x - \tan x}{x^2}$ or equivalent</p> | <p>M1</p> <p>A1 [2]</p> |

Q3.

- 5 (i) Use at least one of $e^{2x} = 9$, $e^y = 2$ and $e^{2y} = 4$ B1
 Obtain given result $58 + 2k = c$ B1 [2]
 AG
- (ii) Differentiate left-hand side term by term, reaching $ae^{2x} + be^y \frac{dy}{dx} + ce^{2y} \frac{dy}{dx}$ M1
 Obtain $12e^{2x} + ke^y \frac{dy}{dx} + 2e^{2y} \frac{dy}{dx}$ A1
 Substitute (ln 3, ln 2) in an attempt involving implicit differentiation at least once, where
 RHS = 0 M1
 Obtain $108 - 12k - 48 = 0$ or equivalent A1
 Obtain $k = 5$ and $c = 68$ A1 [5]

Q4.

- 2 Use correct quotient or product rule M1
 Obtain correct derivative in any form, e.g. $-\frac{3 \ln x}{x^4} + \frac{1}{x^4}$ A1
 Equate derivative to zero and solve for x an equation of the form $\ln x = a$, where $a > 0$ M1
 Obtain answer $\exp(\frac{1}{3})$, or 1.40, from correct work A1 [4]

Q5.

- 6 (i) Obtain $2y \frac{dy}{dx}$ as derivative of y^2 B1
 Obtain $-4y - 4x \frac{dy}{dx}$ as derivative of $-4xy$ B1
 Substitute $x = 2$ and $y = -3$ and find value of $\frac{dy}{dx}$
 (dependent on at least one B1 being earned and $\frac{d(45)}{dx} = 0$) M1
 Obtain $\frac{12}{7}$ or equivalent A1 [4]
- (ii) Substitute $\frac{dy}{dx} = 1$ in an expression involving $\frac{dy}{dx}$, x and y and obtain $ay = bx$ M1
 Obtain $y = x$ or equivalent A1
 Uses $y = x$ in original equation and demonstrate contradiction A1 [3]

Q6.

- 3 Obtain $\frac{dx}{d\theta} = 2 \cos 2\theta - 1$ or $\frac{dy}{d\theta} = -2 \sin 2\theta + 2 \cos \theta$, or equivalent B1
- Use $\frac{dy}{dx} = \frac{dy}{d\theta} \div \frac{dx}{d\theta}$ M1
- Obtain $\frac{dy}{dx} = \frac{-2 \sin 2\theta + 2 \cos \theta}{2 \cos 2\theta - 1}$, or equivalent A1
- At any stage use correct double angle formulae throughout M1
- Obtain the given answer following full and correct working A1 [5]

Q7.

- 4 (i) Use correct quotient or product rule M1
- Obtain correct derivative in any form, e.g. $\frac{2e^{2x}}{x^3} - \frac{3e^{2x}}{x^4}$ A1
- Equate derivative to zero and solve a 2-term equation for non-zero x M1
- Obtain $x = \frac{3}{2}$ correctly A1 [4]
- (ii) Carry out a method for determining the nature of a stationary point, e.g. test derivative either side M1
- Show point is a minimum with no errors seen A1 [2]

Q8.

- 5 (i) Use correct quotient rule or equivalent M1
- Obtain $\frac{(1+e^{2x})2x - (1+x^2)2e^{2x}}{(1+e^{2x})^2}$ or equivalent A1
- Substitute $x = 0$ and obtain $-\frac{1}{2}$ or equivalent A1 [3]
- (ii) Differentiate y^3 and obtain $3y^2 \frac{dy}{dx}$ B1
- Differentiate $5xy$ and obtain $5y + 5x \frac{dy}{dx}$ B1
- Obtain $6x^2 + 5y + 5x \frac{dy}{dx} + 3y^2 \frac{dy}{dx} = 0$ B1

© Cambridge International Examinations 2013

Page 5	Mark Scheme	Syllabus	Paper
	GCE AS/A LEVEL – May/June 2013	9709	31

- Substitute $x = 0, y = 2$ to obtain $-\frac{5}{6}$ or equivalent following correct work B1 [4]

Q9.

- | | | |
|---|--|----------|
| 4 | Use product or quotient rule | M1 |
| | Obtain derivative in any correct form | A1 |
| | Equate derivative to zero and obtain an equation of the form $a \sin 2x = b$, or a quadratic in $\tan x$, $\sin^2 x$, or $\cos^2 x$ | M1* |
| | Carry out correct method for finding one angle | M1(dep*) |
| | Obtain answer, e.g. 0.365 | A1 |
| | Obtain second answer 1.206 and no others in the range (allow 1.21) | A1 [6] |
| | [Ignore answers outside the given range.] | |
| | [Treat answers in degrees, 20.9° and 69.1°, as a misread.] | |

Q10.

- | | | |
|---|---|------------|
| 2 | Use of correct quotient or product rule to differentiate x or t | M1 |
| | Obtain correct $\frac{3}{(2t+3)^2}$ or unsimplified equivalent | A1 |
| | Obtain $-2e^{-2t}$ for derivative of y | B1 |
| | Use $\frac{dy}{dx} = \frac{dy}{dt} \div \frac{dx}{dt}$ or equivalent | M1 |
| | Obtain -6 | cwo A1 [5] |
| | <i>Alternative:</i> | |
| | Eliminate parameter and attempt differentiation $\left(y = e^{\frac{-6x}{1-2x}} \right)$ | B1 |
| | Use correct quotient or product rule | M1 |
| | Use chain rule | M1 |
| | Obtain $\frac{dy}{dx} = \frac{-6}{(1-2x)^2} e^{\frac{-6x}{1-2x}}$ | A1 |
| | Obtain -6 | cwo A1 |

Q11.

- | | | |
|---|--|--------|
| 2 | EITHER: Use chain rule | M1 |
| | obtain $\frac{dx}{dt} = 6 \sin t \cos t$, or equivalent | A1 |
| | obtain $\frac{dy}{dt} = -6 \cos^2 t \sin t$, or equivalent | A1 |
| | Use $\frac{dy}{dx} = \frac{dy}{dt} \div \frac{dx}{dt}$ | M1 |
| | Obtain final answer $\frac{dy}{dx} = -\cos t$ | A1 |
| | OR: Express y in terms of x and use chain rule | M1 |
| | Obtain $\frac{dy}{dx} = k\left(2 - \frac{x}{3}\right)^{\frac{1}{2}}$, or equivalent | A1 |
| | Obtain $\frac{dy}{dx} = -\left(2 - \frac{x}{3}\right)^{\frac{1}{2}}$, or equivalent | A1 |
| | Express derivative in terms of t | M1 |
| | Obtain final answer $\frac{dy}{dx} = -\cos t$ | A1 [5] |

Q12.

2	Use correct quotient or product rule or equivalent Obtain $\frac{(1+e^{2x}).2e^{2x} - e^{2x}.2e^{2x}}{(1+e^{2x})^2}$ or equivalent Substitute $x = \ln 3$ into attempt at first derivative and show use of relevant logarithm property at least once in a correct context Confirm given answer $\frac{9}{50}$ legitimately	M1 A1 M1 A1	[4]
----------	---	----------------------------------	-----

Q13.

8	(i) Differentiate y to obtain $3\sin^2 t \cos t - 3\cos^2 t \sin t$ o.e. Use $\frac{dy}{dx} = \frac{dy}{dt} / \frac{dt}{dx}$ Obtain given result $-3\sin t \cos t$	B1 M1	
	(ii) Identify parameter at origin as $t = \frac{3}{4}\pi$ Use $t = \frac{3}{4}\pi$ to obtain $\frac{3}{2}$	A1cwo B1 B1	[3] [2]
	(iii) Rewrite equation as equation in one trig variable e.g. $\sin 2t = -\frac{2}{3}$, $9 \sin^4 x - 9 \sin^2 x + 1 = 0$, $\tan^2 x + 3 \tan x + 1 = 0$ Find at least one value of t from equation of form $\sin 2t = k$ o.e. Obtain 1.9 Obtain 2.8 and no others	B1 M1 A1 A1	[4]

Q14.

- 7 (i) *EITHER*: State or imply $\frac{1}{x} + \frac{1}{y} \frac{dy}{dx}$ as derivative of $\ln xy$, or equivalent B1
 State or imply $3y^2 \frac{dy}{dx}$ as derivative of y^3 , or equivalent B1
 Equate derivative of LHS to zero and solve for $\frac{dy}{dx}$ M1
 Obtain the given answer A1
- OR* Obtain $xy = \exp(1 + y^3)$ and state or imply $y + x \frac{dy}{dx}$ as derivative of xy B1
 State or imply $3y^2 \frac{dy}{dx} \exp(1 + y^3)$ as derivative of $(1 + y^3)$ B1
 Equate derivatives and solve for $\frac{dy}{dx}$ M1
 Obtain the given answer A1 [4]
 [The M1 is dependent on at least one of the B marks being earned]
- (ii) Equate denominator to zero and solve for y M1*
 Obtain $y = 0.693$ only A1
 Substitute found value in the equation and solve for x M1(dep*)
 Obtain $x = 5.47$ only A1 [4]

Q15.

- 3 (i) Either Use correct quotient rule or equivalent to obtain
- $\frac{dx}{dt} = \frac{4(2t+3) - 8t}{(2t+3)^2}$ or equivalent B1
 Obtain $\frac{dy}{dt} = \frac{4}{2t+3}$ or equivalent B1
 Use $\frac{dy}{dx} = \frac{\frac{dy}{dt}}{\frac{dx}{dt}}$ or equivalent M1
 Obtain $\frac{1}{3}(2t+3)$ or similarly simplified equivalent A1
- Or Express t in terms of x or y e.g. $t = \frac{3x}{4-2x}$ B1
 Obtain Cartesian equation e.g. $y = 2 \ln\left(\frac{6}{2-x}\right)$ B1
 Differentiate and obtain $\frac{dy}{dx} = \frac{2}{2-x}$ M1
 Obtain $\frac{1}{3}(2t+3)$ or similarly simplified equivalent A1 [4]
- (ii) Obtain $2t = 3$ or $t = \frac{3}{2}$ B1
 Substitute in expression for $\frac{dy}{dx}$ and obtain 2 B1 [2]

Q16.

- 1 Use correct quotient or product rule M1
 Obtain correct derivative in any form A1
 Justify the given statement A1 [3]

Q17.

- 4 Use correct product or quotient rule at least once M1*
- Obtain $\frac{dx}{dt} = e^{-t} \sin t - e^{-t} \cos t$ or $\frac{dy}{dt} = e^{-t} \cos t - e^{-t} \sin t$, or equivalent A1
- Use $\frac{dy}{dx} = \frac{dy}{dt} \div \frac{dx}{dt}$ M1
- Obtain $\frac{dy}{dx} = \frac{\sin t - \cos t}{\sin t + \cos t}$, or equivalent A1
- EITHER:* Express $\frac{dy}{dx}$ in terms of $\tan t$ only M1(dep*)
- Show expression is identical to $\tan\left(t - \frac{1}{4}\pi\right)$ A1
- OR:* Express $\tan\left(t - \frac{1}{4}\pi\right)$ in terms of $\tan t$ M1
- Show expression is identical to $\frac{dy}{dx}$ A1 [6]

Q18.

- 4 Differentiate y^3 to obtain $3y^2 \frac{dy}{dx}$ B1
- Use correct product rule at least once *M1
- Obtain $6e^{2x}y + 3e^{2x} \frac{dy}{dx} + e^x y^3 + 3e^x y^2 \frac{dy}{dx}$ as derivative of LHS A1
- Equate derivative of LHS to zero, substitute $x = 0$ and $y = 2$ and find value of $\frac{dy}{dx}$ M1(d*M)
- Obtain $-\frac{4}{3}$ or equivalent as **final answer** A1 [5]

Q19.

- 3 Obtain $\frac{2}{2t+3}$ for derivative of x B1
- Use quotient of product rule, or equivalent, for derivative of y M1
- Obtain $\frac{5}{(2t+3)^2}$ or unsimplified equivalent A1
- Obtain $t = -1$ B1
- Use $\frac{dy}{dx} = \frac{dy}{dt} \div \frac{dx}{dt}$ in algebraic or numerical form M1
- Obtain gradient $\frac{5}{2}$ A1 [6]

Q20.

- 10 (i) Use of product or quotient rule M1
- Obtain $-5e^{-\frac{1}{2}x} \sin 4x + 40e^{-\frac{1}{2}x} \cos 4x$ A1
- Equate $\frac{dy}{dx}$ to zero and obtain $\tan 4x = k$ or $R \cos(4x \pm \alpha)$ M1
- Obtain $\tan 4x = 8$ or $\sqrt{65} \cos\left(4x \pm \tan^{-1} \frac{1}{8}\right)$ A1
- Obtain 0.362 or 20.7° A1
- Obtain 1.147 or 65.7° A1 [6]
- (ii) State or imply that x -coordinates of T_n are increasing by $\frac{1}{4}\pi$ or 45° B1
- Attempt solution of inequality (or equation) of form $x_1 + (n-1)k\pi \leq 25$ M1
- Obtain $n > \frac{4}{\pi}(25 - 0.362) + 1$, following through on their value of x_1 A1✓
- $n = 33$ A1 [4]

Q21.

- 6 Obtain correct derivative of RHS in any form B1
- Obtain correct derivative of LHS in any form B1
- Set $\frac{dy}{dx}$ equal to zero and obtain a horizontal equation M1
- Obtain a correct equation, e.g. $x^2 + y^2 = 1$, from correct work A1
- By substitution in the curve equation, or otherwise, obtain an equation in x^2 or y^2 M1
- Obtain $x = \frac{1}{2}\sqrt{3}$ A1
- Obtain $y = \frac{1}{2}$ A1 7

Q22.

- 4 (i) Use chain rule correctly at least once M1
- Obtain either $\frac{dx}{dt} = \frac{3\sin t}{\cos^4 t}$ or $\frac{dy}{dt} = 3\tan^2 t \sec^2 t$, or equivalent A1
- Use $\frac{dy}{dx} = \frac{dy}{dt} \div \frac{dx}{dt}$ M1
- Obtain the given answer A1 [4]
- (ii) State a correct equation for the tangent in any form B1
- Use Pythagoras M1
- Obtain the given answer A1 [3]

Q23.

- 2 Use correct product rule or correct chain rule to differentiate y M1
- Use $\frac{dy}{dx} = \frac{\frac{dy}{d\theta}}{\frac{dx}{d\theta}}$ M*1
- Obtain $\frac{-4 \cos \theta \sin^2 \theta + 2 \cos^3 \theta}{\sec^2 \theta}$ or equivalent A1
- Express $\frac{dy}{dx}$ in terms of $\cos \theta$ DM*1
- Confirm given answer $6 \cos^5 \theta - 4 \cos^3 \theta$ legitimately A1 [5]

[Online Classes : Megalecture@gmail.com](mailto:Megalecture@gmail.com)
www.youtube.com/megalecture
www.megalecture.com

[Online Classes : Megalecture@gmail.com](mailto:Megalecture@gmail.com)
www.youtube.com/megalecture
www.megalecture.com

[Online Classes : Megalecture@gmail.com](mailto:Megalecture@gmail.com)
www.youtube.com/megalecture
www.megalecture.com