

Q1.

5	The	function	f is such	that $f(x)$	$1 = 2 \sin^2 x$	$-3\cos^2 x$	for $0 \le x \le \pi$.

(i) Express
$$f(x)$$
 in the form $a + b\cos^2 x$, stating the values of a and b.

(ii) State the greatest and least values of
$$f(x)$$
. [2]

[2]

(iii) Solve the equation
$$f(x) + 1 = 0$$
. [3]

Q2.

- 9 The function f is defined by $f: x \mapsto 2x^2 12x + 7$ for $x \in \mathbb{R}$.
 - (i) Express f(x) in the form $a(x-b)^2 c$. [3]
 - (ii) State the range of f. [1]
 - (iii) Find the set of values of x for which f(x) < 21. [3]

The function g is defined by $g: x \mapsto 2x + k$ for $x \in \mathbb{R}$.

(iv) Find the value of the constant k for which the equation gf(x) = 0 has two equal roots. [4]

Q3.

- 3 The function $f: x \mapsto a + b \cos x$ is defined for $0 \le x \le 2\pi$. Given that f(0) = 10 and that $f(\frac{2}{3}\pi) = 1$, find
 - (i) the values of a and b, [2]
 - (ii) the range of f, [1]
 - (iii) the exact value of $f(\frac{5}{6}\pi)$. [2]

Q4.

- 10 The function $f: x \mapsto 2x^2 8x + 14$ is defined for $x \in \mathbb{R}$.
 - (i) Find the values of the constant k for which the line y + kx = 12 is a tangent to the curve y = f(x). [4]
 - (ii) Express f(x) in the form $a(x+b)^2 + c$, where a, b and c are constants. [3]
 - (iii) Find the range of f. [1]

The function $g: x \mapsto 2x^2 - 8x + 14$ is defined for $x \ge A$.

- (iv) Find the smallest value of A for which g has an inverse. [1]
- (v) For this value of A, find an expression for $g^{-1}(x)$ in terms of x. [3]

Q5.



11 Functions f and g are defined for $x \in \mathbb{R}$ by

$$f: x \mapsto 2x + 1,$$

 $g: x \mapsto x^2 - 2.$

- (i) Find and simplify expressions for fg(x) and gf(x). [2]
- (ii) Hence find the value of a for which fg(a) = gf(a). [3]
- (iii) Find the value of b ($b \neq a$) for which g(b) = b. [2]
- (iv) Find and simplify an expression for $f^{-1}g(x)$. [2]

The function h is defined by

$$h: x \mapsto x^2 - 2$$
, for $x \le 0$.

(v) Find an expression for $h^{-1}(x)$. [2]

Q6.

10 Functions f and g are defined by

f:
$$x \mapsto 3x - 4$$
, $x \in \mathbb{R}$,
g: $x \mapsto 2(x - 1)^3 + 8$, $x > 1$.

- (i) Evaluate fg(2). [2]
- (ii) Sketch in a single diagram the graphs of y = f(x) and $y = f^{-1}(x)$, making clear the relationship between the graphs. [3]
- (iii) Obtain an expression for g'(x) and use your answer to explain why g has an inverse. [3]
- (iv) Express each of $f^{-1}(x)$ and $g^{-1}(x)$ in terms of x. [4]

Q7.

- 8 The function $f: x \mapsto x^2 4x + k$ is defined for the domain $x \ge p$, where k and p are constants.
 - (i) Express f(x) in the form $(x+a)^2 + b + k$, where a and b are constants. [2]
 - (ii) State the range of f in terms of k. [1]
 - (iii) State the smallest value of p for which f is one-one. [1]
 - (iv) For the value of p found in part (iii), find an expression for $f^{-1}(x)$ and state the domain of f^{-1} , giving your answers in terms of k. [4]

Q8.



- 11 The function f is such that $f(x) = 8 (x-2)^2$, for $x \in \mathbb{R}$.
 - (i) Find the coordinates and the nature of the stationary point on the curve y = f(x). [3]

The function g is such that $g(x) = 8 - (x - 2)^2$, for $k \le x \le 4$, where k is a constant.

(ii) State the smallest value of k for which g has an inverse. [1]

For this value of k,

- (iii) find an expression for $g^{-1}(x)$, [3]
- (iv) sketch, on the same diagram, the graphs of y = g(x) and $y = g^{-1}(x)$. [3]

Q9.

- 8 (i) Express $2x^2 12x + 13$ in the form $a(x+b)^2 + c$, where a, b and c are constant. [3]
 - (ii) The function f is defined by $f(x) = 2x^2 12x + 13$ for $x \ge k$, where k is a constant. It is given that f is a one-one function. State the smallest possible value of k. [1]

The value of k is now given to be 7.

- (iii) Find the range of f. [1]
- (iv) Find an expression for $f^{-1}(x)$ and state the domain of f. [5]

Q10.

- 10 The function f is defined by $f: x \mapsto 2x + k$, $x \in \mathbb{R}$, where k is a constant.
 - (i) In the case where k = 3, solve the equation ff(x) = 25. [2]

The function g is defined by $g: x \mapsto x^2 - 6x + 8, x \in \mathbb{R}$.

(ii) Find the set of values of k for which the equation f(x) = g(x) has no real solutions. [3]

The function h is defined by h: $x \mapsto x^2 - 6x + 8, x > 3$.

(iii) Find an expression for $h^{-1}(x)$. [4]

Q11.



10 Functions f and g are defined by

$$f: x \mapsto 2x + 1, \quad x \in \mathbb{R}, \quad x > 0,$$
$$g: x \mapsto \frac{2x - 1}{x + 3}, \quad x \in \mathbb{R}, \quad x \neq -3.$$

(i) Solve the equation gf(x) = x.

[3]

(ii) Express $f^{-1}(x)$ and $g^{-1}(x)$ in terms of x.

[4]

(iii) Show that the equation $g^{-1}(x) = x$ has no solutions.

- [3]
- (iv) Sketch in a single diagram the graphs of y = f(x) and $y = f^{-1}(x)$, making clear the relationship between the graphs. [3]

Q12.

3 Functions f and g are defined for $x \in \mathbb{R}$ by

$$f: x \mapsto 2x + 3$$
,
 $g: x \mapsto x^2 - 2x$.

Express gf(x) in the form $a(x+b)^2 + c$, where a, b and c are constants.

[5]

Q13.

- 6 A curve has equation y = f(x). It is given that $f'(x) = 3x^2 + 2x 5$.
 - (i) Find the set of values of x for which f is an increasing function.

[3]


(ii) Given that the curve passes through (1, 3), find f(x).

[4]

Q14.



7



The diagram shows the function f defined for $0 \le x \le 6$ by

$$x \mapsto \frac{1}{2}x^2$$
 for $0 \le x \le 2$,
 $x \mapsto \frac{1}{2}x + 1$ for $2 < x \le 6$.

(i) State the range of f.

[1]

(ii) Copy the diagram and on your copy sketch the graph of $y = f^{-1}(x)$.

- [2]
- (iii) Obtain expressions to define $f^{-1}(x)$, giving the set of values of x for which each expression is valid. [4]

Q15.

11 Functions f and g are defined by

f
$$x = 2x^2 - 8x + 10$$
 for $0 \le x \le 2$,
g $x \mapsto x$ for $0 \le x \le 10$

- (i) Express f(x) in the form $a(x+b)^2 + c$, where a, b and c are constants.
- [3]

(ii) State the range of f.

[1]

(iii) State the domain of f^{-1} .

- [1]
- (iv) Sketch on the same diagram the graphs of y = f(x), y = g(x) and $y = f^{-1}(x)$, making clear the relationship between the graphs. [4]
- (v) Find an expression for $f^{-1}(x)$.

[3]

Q16.



9 Functions f and g are defined by

f:
$$x \mapsto 2x + 3$$
 for $x \le 0$,
g: $x \mapsto x^2 - 6x$ for $x \le 3$.

- (i) Express $f^{-1}(x)$ in terms of x and solve the equation $f(x) = f^{-1}(x)$. [3]
- (ii) On the same diagram sketch the graphs of y = f(x) and $y = f^{-1}(x)$, showing the coordinates of their point of intersection and the relationship between the graphs. [3]
- (iii) Find the set of values of x which satisfy $gf(x) \le 16$. [5]

Q17.

- 10 The function f is defined by $f(x) = 4x^2 24x + 11$, for $x \in \mathbb{R}$.
 - (i) Express f(x) in the form a(x b)² + c and hence state the coordinates of the vertex of the graph of y = f(x).

The function g is defined by $g(x) = 4x^2 - 24x + 11$, for $x \le 1$.

- (ii) State the range of g. [2]
- (iii) Find an expression for $g^{-1}(x)$ and state the domain of g^{-1} . [4]

Q18.

6 The functions f and g are defined for $-\frac{1}{2}\pi \leqslant x \leqslant \frac{1}{2}\pi$ by

$$f(x) = \frac{1}{2}x + \frac{1}{6}\pi,$$

$$g(x) = \cos x.$$


Solve the following equations for $-\frac{1}{2}\pi \le x \le \frac{1}{2}\pi$.

- (i) gf (x) = 1, giving your answer in terms of π . [2]
- (ii) fg(x) = 1, giving your answers correct to 2 decimal places. [4]

Q19.



7



- (i) The diagram shows part of the curve $y = 11 x^2$ and part of the straight line y = 5 x meeting at the point A(p, q), where p and q are positive constants. Find the values of p and q. [3]
- (ii) The function f is defined for the domain $x \ge 0$ by

$$f(x) = \begin{cases} 11 - x^2 & \text{for } 0 \le x \le p, \\ 5 - x & \text{for } x > p. \end{cases}$$

Express $f^{-1}(x)$ in a similar way.

[5]

Q20.

5 The function f is defined by

 $f: x \rightarrow x^2 + 1 \text{ for } x \ge 0.$

(i) Define in a similar way the inverse function f⁻¹

(ii) Solve the equation $f(x) = \frac{185}{16}$.

[3]

[3]

Q21.

- 10 The function f is defined by $f: x \mapsto x^2 + 4x$ for $x \ge c$, where c is a constant. It is given that f is a one-one function.
 - (i) State the range of f in terms of c and find the smallest possible value of c. [3]

The function g is defined by $g: x \mapsto ax + b$ for $x \ge 0$, where a and b are positive constants. It is given that, when c = 0, gf(1) = 11 and fg(1) = 21.

(ii) Write down two equations in a and b and solve them to find the values of a and b. [6]

