


These are P2 questions(all variants) as the syllabus is same as P3 :)

Q1.

3



The diagram shows the curve $y = e^{2x}$. The shaded region R is bounded by the curve and by the lines $x = 0$, $y = 0$ and $x = p$.

- (i) Find, in terms of p , the area of R . [3]
- (ii) Hence calculate the value of p for which the area of R is equal to 5. Give your answer correct to 2 significant figures. [3]

Q2.

- 7 (i) By expanding $\cos(2x + x)$, show that

$$\cos 3x = 4 \cos^3 x - 3 \cos x. \quad [5]$$

- (ii) Hence, or otherwise, show that

$$\int_0^{\frac{1}{2}\pi} \cos^3 x \, dx = \frac{2}{3}. \quad [5]$$

Q3.

- 7 (i) By expanding $\sin(2x + x)$ and using double-angle formulae, show that

$$\sin 3x = 3 \sin x - 4 \sin^3 x. \quad [5]$$

- (ii) Hence show that

$$\int_0^{\frac{1}{3}\pi} \sin^3 x \, dx = \frac{5}{24}. \quad [5]$$

Q4.

6 (i) Express $\cos^2 x$ in terms of $\cos 2x$. [1]

(ii) Hence show that

$$\int_0^{\frac{1}{3}\pi} \cos^2 x \, dx = \frac{1}{6}\pi + \frac{1}{8}\sqrt{3}. \quad [4]$$

(iii) By using an appropriate trigonometrical identity, deduce the exact value of


$$\int_0^{\frac{1}{3}\pi} \sin^2 x \, dx. \quad [3]$$

Q5.

3 Find the exact value of $\int_0^{\frac{1}{6}\pi} (\cos 2x + \sin x) \, dx$. [5]

Q6.

3



The diagram shows the curve $y = \frac{1}{1 + \sqrt{x}}$ for values of x from 0 to 2.

(i) Use the trapezium rule with two intervals to estimate the value of

$$\int_0^2 \frac{1}{1 + \sqrt{x}} \, dx,$$

giving your answer correct to 2 decimal places. [3]


(ii) State, with a reason, whether the trapezium rule gives an under-estimate or an over-estimate of the true value of the integral in part (i). [1]

Q7.

2 Show that $\int_0^6 \frac{1}{x+2} dx = 2 \ln 2$. [4]

Q8.

2



The diagram shows part of the curve $y = xe^{-x}$. The shaded region R is bounded by the curve and by the lines $x = 2$, $x = 3$ and $y = 0$.

- (i) Use the trapezium rule with two intervals to estimate the area of R , giving your answer correct to 2 decimal places. [3]
- (ii) State, with a reason, whether the trapezium rule gives an under-estimate or an over-estimate of the true value of the area of R . [1]

Q9.

4 (a) Show that $\int_0^{\frac{1}{2}\pi} \cos 2x dx = \frac{1}{2}$. [2]

(b) By using an appropriate trigonometrical identity, find the exact value of

$$\int_{\frac{1}{6}\pi}^{\frac{1}{3}\pi} 3 \tan^2 x dx. \quad [4]$$


Q10.

6 (a) Find $\int 4e^x(3 + e^{2x}) dx$. [4]

(b) Show that $\int_{-\frac{1}{4}\pi}^{\frac{1}{4}\pi} (3 + 2 \tan^2 \theta) d\theta = \frac{1}{2}(8 + \pi)$. [4]

Q11.

2



The diagram shows the curve $y = \sqrt{1+x^3}$. Region A is bounded by the curve and the lines $x = 0$, $x = 2$ and $y = 0$. Region B is bounded by the curve and the lines $x = 0$ and $y = 3$.

- (i) Use the trapezium rule with two intervals to find an approximation to the area of region A. Give your answer correct to 2 decimal places. [3]
- (ii) Deduce an approximation to the area of region B and explain why this approximation underestimates the true area of region B. [2]

Q12.

- 4 (a) Find the value of $\int_0^{\pi} \sin\left(\frac{1}{2}x\right) dx$. [3]
- (b) Find $\int e^{-x}(1 + e^x) dx$. [3]

Q13.


- 7 (i) Show that $(2 \sin x + \cos x)^2$ can be written in the form $\frac{5}{2} + 2 \sin 2x - \frac{3}{2} \cos 2x$. [5]
- (ii) Hence find the exact value of $\int_0^{\frac{1}{4}\pi} (2 \sin x + \cos x)^2 dx$. [4]

Q14.

- 7 (i) Show that $\tan^2 x + \cos^2 x \equiv \sec^2 x + \frac{1}{2} \cos 2x - \frac{1}{2}$ and hence find the exact value of

$$\int_0^{\frac{1}{4}\pi} (\tan^2 x + \cos^2 x) dx. \quad [7]$$

(ii)



The region enclosed by the curve $y = \tan x + \cos x$ and the lines $x = 0$, $x = \frac{1}{4}\pi$ and $y = 0$ is shown in the diagram. Find the exact volume of the solid produced when this region is rotated completely about the x -axis. [4]

Q15.

- 3 (i) Show that $12 \sin^2 x \cos^2 x \equiv \frac{3}{2}(1 - \cos 4x)$. [3]

(ii) Hence show that

$$\int_{\frac{1}{4}\pi}^{\frac{3}{4}\pi} 12 \sin^2 x \cos^2 x dx = \frac{\pi}{8} + \frac{3\sqrt{3}}{16}. \quad [3]$$


Q16.

- 1 A curve is such that $\frac{dy}{dx} = \frac{4}{7-2x}$. The point (3, 2) lies on the curve. Find the equation of the curve. [4]

Q17.

6 (a) Find the value of $\int_0^{\frac{1}{2}\pi} (\sin 2x + \cos x) dx$. [4]

(b)



The diagram shows part of the curve $y = \frac{1}{x+1}$. The shaded region R is bounded by the curve and by the lines $x = 1$, $y = 0$ and $x = p$.

(i) Find, in terms of p , the area of R . [3]

(ii) Hence find, correct to 1 decimal place, the value of p for which the area of R is equal to 2. [2]

Q18.

7 (i) Given that $y = \tan 2x$, find $\frac{dy}{dx}$. [2]

(ii) Hence, or otherwise, show that

$$\int_0^{\frac{1}{8}\pi} \sec^2 2x dx = \frac{1}{2}\sqrt{3},$$

and, by using an appropriate trigonometrical identity, find the exact value of $\int_0^{\frac{1}{8}\pi} \tan^2 2x dx$. [6]

(iii) Use the identity $\cos 4x \equiv 2 \cos^2 2x - 1$ to find the exact value of

$$\int_0^{\frac{1}{8}\pi} \frac{1}{1 + \cos 4x} dx. [2]$$

Q19.

1 Show that

$$\int_1^4 \frac{1}{2x+1} dx = \frac{1}{2} \ln 3. [4]$$

Q20.

- 7 (i) Prove the identity

$$(\cos x + 3 \sin x)^2 \equiv 5 - 4 \cos 2x + 3 \sin 2x. \quad [4]$$

- (ii) Using the identity, or otherwise, find the exact value of

$$\int_0^{\frac{1}{4}\pi} (\cos x + 3 \sin x)^2 dx. \quad [4]$$

Q21.

5 Show that $\int_1^2 \left(\frac{1}{x} - \frac{4}{2x+1} \right) dx = \ln \frac{18}{25}.$ [6]

Q22.

- 5 (i) Express $\cos^2 2x$ in terms of $\cos 4x.$ [2]

(ii) Hence find the exact value of $\int_0^{\frac{1}{8}\pi} \cos^2 2x dx.$ [4]

Q23.

- 3 (i) Use the trapezium rule with two intervals to estimate the value of

$$\int_0^{\frac{1}{3}\pi} \sec x dx,$$

giving your answer correct to 2 decimal places. [3]

- (ii) Using a sketch of the graph of $y = \sec x$ for $0 \leq x \leq \frac{1}{3}\pi$, explain whether the trapezium rule gives an under-estimate or an over-estimate of the true value of the integral in part (i). [2]

Q24.

8 (a) Find the exact value of $\int_0^{\frac{1}{3}\pi} (\sin 2x + \sec^2 x) dx.$ [5]

(b) Show that $\int_1^4 \left(\frac{1}{2x} + \frac{1}{x+1} \right) dx = \ln 5.$ [4]

Q25.

3 Show that $\int_0^1 (e^x + 1)^2 dx = \frac{1}{2}e^2 + 2e - \frac{3}{2}$. [5]

Q26.

4 (a) Find $\int e^{1-2x} dx$. [2]

(b) Express $\sin^2 3x$ in terms of $\cos 6x$ and hence find $\int \sin^2 3x dx$. [4]

Q27.

2 Show that $\int_2^6 \frac{2}{4x+1} dx = \ln \frac{5}{3}$. [5]

Q28.

8 (i) By first expanding $\cos(2x + x)$, show that

$$\cos 3x = 4 \cos^3 x - 3 \cos x. \quad [5]$$

(ii) Hence show that

$$\int_0^{\frac{1}{6}\pi} (2 \cos^3 x - \cos x) dx = \frac{5}{12}. \quad [5]$$

Q29.

4 Find the exact value of the positive constant k for which

$$\int_0^k e^{4x} dx = \int_0^{2k} e^x dx. \quad [6]$$

Q30.

4 (i) Express $\cos^2 x$ in terms of $\cos 2x$. [1]

(ii) Hence show that

$$\int_0^{\frac{1}{6}\pi} (\cos^2 x + \sin 2x) dx = \frac{1}{8}\sqrt{3} + \frac{1}{12}\pi + \frac{1}{4}. \quad [5]$$

Q31.

- 6 (a) Use the trapezium rule with two intervals to estimate the value of

$$\int_0^1 \frac{1}{6 + 2e^x} dx,$$

giving your answer correct to 2 decimal places. [3]

(b) Find $\int \frac{(e^x - 2)^2}{e^{2x}} dx$. [4]

Q32.

6 (a) Find $\int 4e^{-\frac{1}{2}x} dx$. [2]

(b) Show that $\int_1^3 \frac{6}{3x-1} dx = \ln 16$. [5]

Q33.

- 6 (a) Find

(i) $\int \frac{e^{2x} + 6}{e^{2x}} dx$, [3]

(ii) $\int 3 \cos^2 x dx$. [3]

- (b) Use the trapezium rule with 2 intervals to estimate the value of

$$\int_1^2 \frac{6}{\ln(x+2)} dx,$$

giving your answer correct to 2 decimal places. [3]

Q34.

1 (i) Find $\int \frac{2}{4x-1} dx$. [2]

(ii) Hence find $\int_1^7 \frac{2}{4x-1} dx$, expressing your answer in the form $\ln a$, where a is an integer. [3]

Q35.

6 (a) Find $\int (\sin x - \cos x)^2 dx$. [4]

(b) (i) Use the trapezium rule with 2 intervals to estimate the value of

$$\int_{\frac{1}{4}\pi}^{\frac{1}{2}\pi} \operatorname{cosec} x \, dx,$$

giving your answer correct to 3 decimal places. [3]

(ii) Using a sketch of the graph of $y = \operatorname{cosec} x$ for $0 < x \leq \frac{1}{2}\pi$, explain whether the trapezium rule gives an under-estimate or an over-estimate of the true value of the integral in part (i). [2]

Q36.

5 (i) Prove that $\tan \theta + \cot \theta \equiv \frac{2}{\sin 2\theta}$. [3]

(ii) Hence

(a) find the exact value of $\tan \frac{1}{8}\pi + \cot \frac{1}{8}\pi$. [2]

(b) evaluate $\int_0^{\frac{1}{2}\pi} \frac{6}{\tan \theta + \cot \theta} d\theta$. [3]

Q37.

6 (a) Show that $\int_6^{16} \frac{6}{2x-7} dx = \ln 125$. [5]

(b) Use the trapezium rule with four intervals to find an approximation to

$$\int_1^{17} \log_{10} x \, dx,$$

giving your answer correct to 3 significant figures. [3]

Q38.

3 (a) Find $\int 4 \cos\left(\frac{1}{3}x + 2\right) dx$. [2]

(b) Use the trapezium rule with three intervals to find an approximation to

$$\int_0^{12} \sqrt{4+x^2} dx,$$

giving your answer correct to 3 significant figures. [3]

Q39.

- 1 Use the trapezium rule with four intervals to find an approximation to

$$\int_1^5 |2^x - 8| dx. \quad [3]$$

Q40.

3 (a) Find $\int 4 \cos^2\left(\frac{1}{2}\theta\right) d\theta$. [3]

(b) Find the exact value of $\int_{-1}^6 \frac{1}{2x+3} dx$. [4]

Q41.

2 (i) Find $\int_0^a (e^{-x} + 6e^{-3x}) dx$, where a is a positive constant. [4]

(ii) Deduce the value of $\int_0^{\infty} (e^{-x} + 6e^{-3x}) dx$. [1]

P3 (variant1 and 3)

Q1.

- 4 (i) Using the expansions of $\cos(3x - x)$ and $\cos(3x + x)$, prove that

$$\frac{1}{2}(\cos 2x - \cos 4x) \equiv \sin 3x \sin x. \quad [3]$$

- (ii) Hence show that

$$\int_{\frac{1}{3}\pi}^{\frac{1}{2}\pi} \sin 3x \sin x dx = \frac{1}{8}\sqrt{3}. \quad [3]$$

Q2.

8 (i) Express $\frac{2}{(x+1)(x+3)}$ in partial fractions. [2]

- (ii) Using your answer to part (i), show that

$$\left(\frac{2}{(x+1)(x+3)}\right)^2 \equiv \frac{1}{(x+1)^2} - \frac{1}{x+1} + \frac{1}{x+3} + \frac{1}{(x+3)^2}. \quad [2]$$

(iii) Hence show that $\int_0^1 \frac{4}{(x+1)^2(x+3)^2} dx = \frac{7}{12} - \ln \frac{3}{2}$. [5]

Q3.

7 (i) Prove the identity $\cos 3\theta \equiv 4 \cos^3 \theta - 3 \cos \theta$. [4]

(ii) Using this result, find the exact value of

$$\int_{\frac{1}{3}\pi}^{\frac{1}{2}\pi} \cos^3 \theta \, d\theta. \quad [4]$$

Q4.

7 The integral I is defined by $I = \int_0^2 4t^3 \ln(t^2 + 1) \, dt$.

(i) Use the substitution $x = t^2 + 1$ to show that $I = \int_1^5 (2x - 2) \ln x \, dx$. [3]

(ii) Hence find the exact value of I . [5]

Q5.

10 The number of birds of a certain species in a forested region is recorded over several years. At time t years, the number of birds is N , where N is treated as a continuous variable. The variation in the number of birds is modelled by

$$\frac{dN}{dt} = \frac{N(1800 - N)}{3600}.$$

It is given that $N = 300$ when $t = 0$.

(i) Find an expression for N in terms of t . [9]

(ii) According to the model, how many birds will there be after a long time? [1]

Q6.

3 Show that $\int_0^1 (1-x)e^{-\frac{1}{2}x} \, dx = 4e^{-\frac{1}{2}} - 2$. [5]

Q7.

- 9 In a chemical reaction, a compound X is formed from two compounds Y and Z . The masses in grams of X , Y and Z present at time t seconds after the start of the reaction are x , $10 - x$ and $20 - x$ respectively. At any time the rate of formation of X is proportional to the product of the masses of Y and Z present at the time. When $t = 0$, $x = 0$ and $\frac{dx}{dt} = 2$.

(i) Show that x and t satisfy the differential equation

$$\frac{dx}{dt} = 0.01(10 - x)(20 - x). \quad [1]$$

(ii) Solve this differential equation and obtain an expression for x in terms of t . [9]

(iii) State what happens to the value of x when t becomes large. [1]

Q8.

- 7 The variables x and y are related by the differential equation

$$\frac{dy}{dx} = \frac{6xe^{3x}}{y^2}.$$

It is given that $y = 2$ when $x = 0$. Solve the differential equation and hence find the value of y when $x = 0.5$, giving your answer correct to 2 decimal places. [8]

Q9.

- 5 Given that $y = 0$ when $x = 1$, solve the differential equation

$$xy \frac{dy}{dx} = y^2 + 4,$$

obtaining an expression for y^2 in terms of x . [6]

Q10.

- 4 Given that $x = 1$ when $t = 0$, solve the differential equation

$$\frac{dx}{dt} = \frac{1}{x} - \frac{x}{4},$$

obtaining an expression for x^2 in terms of t . [7]

Q11.

- 9 By first expressing $\frac{4x^2 + 5x + 3}{2x^2 + 5x + 2}$ in partial fractions, show that

$$\int_0^4 \frac{4x^2 + 5x + 3}{2x^2 + 5x + 2} dx = 8 - \ln 9. \quad [10]$$

Q12.

- 5 In a certain chemical process a substance A reacts with another substance B . The masses in grams of A and B present at time t seconds after the start of the process are x and y respectively. It is given that $\frac{dy}{dt} = -0.6xy$ and $x = 5e^{-3t}$. When $t = 0$, $y = 70$.
- (i) Form a differential equation in y and t . Solve this differential equation and obtain an expression for y in terms of t . [6]
- (ii) The percentage of the initial mass of B remaining at time t is denoted by p . Find the exact value approached by p as t becomes large. [2]

Q13.

- 8 Let $f(x) = \frac{4x^2 - 7x - 1}{(x+1)(2x-3)}$.
- (i) Express $f(x)$ in partial fractions. [5]
- (ii) Show that $\int_2^6 f(x) dx = 8 - \ln\left(\frac{49}{3}\right)$. [5]

Q14.

- 8 (a) Show that $\int_2^4 4x \ln x dx = 56 \ln 2 - 12$. [5]
- (b) Use the substitution $u = \sin 4x$ to find the exact value of $\int_0^{\frac{1}{24}\pi} \cos^3 4x dx$. [5]

Q15.

- 10 Liquid is flowing into a small tank which has a leak. Initially the tank is empty and, t minutes later, the volume of liquid in the tank is $V \text{ cm}^3$. The liquid is flowing into the tank at a constant rate of 80 cm^3 per minute. Because of the leak, liquid is being lost from the tank at a rate which, at any instant, is equal to $kV \text{ cm}^3$ per minute where k is a positive constant.

(i) Write down a differential equation describing this situation and solve it to show that

$$V = \frac{1}{k}(80 - 80e^{-kt}). \quad [7]$$

(ii) It is observed that $V = 500$ when $t = 15$, so that k satisfies the equation

$$k = \frac{4 - 4e^{-15k}}{25}.$$

Use an iterative formula, based on this equation, to find the value of k correct to 2 significant figures. Use an initial value of $k = 0.1$ and show the result of each iteration to 4 significant figures. [3]

(iii) Determine how much liquid there is in the tank 20 minutes after the liquid started flowing, and state what happens to the volume of liquid in the tank after a long time. [2]

Q16.

- 4 (i) Express $(\sqrt{3}) \cos x + \sin x$ in the form $R \cos(x - \alpha)$, where $R > 0$ and $0 < \alpha < \frac{1}{2}\pi$, giving the exact values of R and α . [3]

(ii) Hence show that

$$\int_{\frac{1}{8}\pi}^{\frac{1}{2}\pi} \frac{1}{((\sqrt{3}) \cos x + \sin x)^2} dx = \frac{1}{4}\sqrt{3}. \quad [4]$$

Q17.

- 8 The variables x and t satisfy the differential equation

$$t \frac{dx}{dt} = \frac{k - x^3}{2x^2},$$

for $t > 0$, where k is a constant. When $t = 1$, $x = 1$ and when $t = 4$, $x = 2$.

(i) Solve the differential equation, finding the value of k and obtaining an expression for x in terms of t . [9]

(ii) State what happens to the value of x as t becomes large. [1]

Q18.

- 5 (i) Prove the identity $\cos 4\theta - 4 \cos 2\theta + 3 \equiv 8 \sin^4 \theta$. [4]
(ii) Using this result find, in simplified form, the exact value of

$$\int_{\frac{1}{6}\pi}^{\frac{1}{3}\pi} \sin^4 \theta \, d\theta. \quad [4]$$

Q19.

- 10 In a model of the expansion of a sphere of radius r cm, it is assumed that, at time t seconds after the start, the rate of increase of the surface area of the sphere is proportional to its volume. When $t = 0$, $r = 5$ and $\frac{dr}{dt} = 2$.

- (i) Show that r satisfies the differential equation

$$\frac{dr}{dt} = 0.08r^2. \quad [4]$$

[The surface area A and volume V of a sphere of radius r are given by the formulae $A = 4\pi r^2$, $V = \frac{4}{3}\pi r^3$.]

- (ii) Solve this differential equation, obtaining an expression for r in terms of t . [5]
(iii) Deduce from your answer to part (ii) the set of values that t can take, according to this model. [1]

Q20.

5 Let $I = \int_0^1 \frac{x^2}{\sqrt{4-x^2}} \, dx$.

- (i) Using the substitution $x = 2 \sin \theta$, show that

$$I = \int_0^{\frac{1}{6}\pi} 4 \sin^2 \theta \, d\theta. \quad [3]$$

- (ii) Hence find the exact value of I . [4]

Q21.

- 10 A certain substance is formed in a chemical reaction. The mass of substance formed t seconds after the start of the reaction is x grams. At any time the rate of formation of the substance is proportional to $(20 - x)$. When $t = 0$, $x = 0$ and $\frac{dx}{dt} = 1$.

(i) Show that x and t satisfy the differential equation

$$\frac{dx}{dt} = 0.05(20 - x). \quad [2]$$

(ii) Find, in any form, the solution of this differential equation. [5]

(iii) Find x when $t = 10$, giving your answer correct to 1 decimal place. [2]

(iv) State what happens to the value of x as t becomes very large. [1]

Q22.

5 Show that $\int_0^7 \frac{2x+7}{(2x+1)(x+2)} dx = \ln 50$. [7]

Q23.

- 9 A biologist is investigating the spread of a weed in a particular region. At time t weeks after the start of the investigation, the area covered by the weed is A m². The biologist claims that the rate of increase of A is proportional to $\sqrt{(2A - 5)}$.

(i) Write down a differential equation representing the biologist's claim. [1]

(ii) At the start of the investigation, the area covered by the weed was 7 m² and, 10 weeks later, the area covered was 27 m². Assuming that the biologist's claim is correct, find the area covered 20 weeks after the start of the investigation. [9]

Q24.

- 4 The variables x and θ are related by the differential equation

$$\sin 2\theta \frac{dx}{d\theta} = (x + 1) \cos 2\theta,$$

where $0 < \theta < \frac{1}{2}\pi$. When $\theta = \frac{1}{12}\pi$, $x = 0$. Solve the differential equation, obtaining an expression for x in terms of θ , and simplifying your answer as far as possible. [7]

Q25.

8 Let $f(x) = \frac{12 + 8x - x^2}{(2 - x)(4 + x^2)}$.

(i) Express $f(x)$ in the form $\frac{A}{2 - x} + \frac{Bx + C}{4 + x^2}$. [4]

(ii) Show that $\int_0^1 f(x) dx = \ln\left(\frac{25}{2}\right)$. [5]

Q26.

- 4 During an experiment, the number of organisms present at time t days is denoted by N , where N is treated as a continuous variable. It is given that

$$\frac{dN}{dt} = 1.2e^{-0.02t}N^{0.5}.$$

When $t = 0$, the number of organisms present is 100.

(i) Find an expression for N in terms of t . [6]

(ii) State what happens to the number of organisms present after a long time. [1]

Q27.

- 10 (i) Use the substitution $u = \tan x$ to show that, for $n \neq -1$,

$$\int_0^{\frac{1}{2}\pi} (\tan^{n+2}x + \tan^n x) dx = \frac{1}{n+1}. \quad [4]$$

(ii) Hence find the exact value of

(a) $\int_0^{\frac{1}{2}\pi} (\sec^4 x - \sec^2 x) dx$, [3]

(b) $\int_0^{\frac{1}{2}\pi} (\tan^9 x + 5 \tan^7 x + 5 \tan^5 x + \tan^3 x) dx$. [3]

Q28.

- 6 The variables x and y are related by the differential equation

$$x \frac{dy}{dx} = 1 - y^2.$$

When $x = 2$, $y = 0$. Solve the differential equation, obtaining an expression for y in terms of x . [8]

Q29.

- 4 The variables x and y are related by the differential equation


$$(x^2 + 4) \frac{dy}{dx} = 6xy.$$

It is given that $y = 32$ when $x = 0$. Find an expression for y in terms of x .

[6]

Q30.

7



The diagram shows part of the curve $y = \sin^3 2x \cos^3 2x$. The shaded region shown is bounded by the curve and the x -axis and its exact area is denoted by A .

- (i) Use the substitution $u = \sin 2x$ in a suitable integral to find the value of A . [6]

- (ii) Given that $\int_0^{k\pi} |\sin^3 2x \cos^3 2x| dx = 40A$, find the value of the constant k . [2]

Q31.

- 3 Find the exact value of $\int_1^4 \frac{\ln x}{\sqrt{x}} dx$. [5]


Q32.

- 5 (i) Prove that $\cot \theta + \tan \theta \equiv 2 \operatorname{cosec} 2\theta$. [3]

- (ii) Hence show that $\int_{\frac{1}{6}\pi}^{\frac{1}{3}\pi} \operatorname{cosec} 2\theta d\theta = \frac{1}{2} \ln 3$. [4]

Q33.

10



A tank containing water is in the form of a cone with vertex C . The axis is vertical and the semi-vertical angle is 60° , as shown in the diagram. At time $t = 0$, the tank is full and the depth of water is H . At this instant, a tap at C is opened and water begins to flow out. The volume of water in the tank decreases at a rate proportional to \sqrt{h} , where h is the depth of water at time t . The tank becomes empty when $t = 60$.

(i) Show that h and t satisfy a differential equation of the form

$$\frac{dh}{dt} = -Ah^{-\frac{3}{2}},$$

where A is a positive constant.

[4]

(ii) Solve the differential equation given in part (i) and obtain an expression for t in terms of h and H .

[6]

(iii) Find the time at which the depth reaches $\frac{1}{2}H$.

[1]

[The volume V of a cone of vertical height h and base radius r is given by $V = \frac{1}{3}\pi r^2 h$.]


Q34.

2 Use the substitution $u = 3x + 1$ to find $\int \frac{3x}{3x+1} dx$.

[4]

Q35.

10



A particular solution of the differential equation

$$3y^2 \frac{dy}{dx} = 4(y^3 + 1) \cos^2 x$$

is such that $y = 2$ when $x = 0$. The diagram shows a sketch of the graph of this solution for $0 \leq x \leq 2\pi$; the graph has stationary points at A and B . Find the y -coordinates of A and B , giving each coordinate correct to 1 decimal place. [10]

Q36.

- 2 Use the substitution $u = 1 + 3 \tan x$ to find the exact value of

$$\int_0^{\frac{1}{4}\pi} \frac{\sqrt{1 + 3 \tan x}}{\cos^2 x} dx. \quad [5]$$

Q37.

- 4 The variables x and y are related by the differential equation

$$\frac{dy}{dx} = \frac{6ye^{3x}}{2 + e^{3x}}.$$

Given that $y = 36$ when $x = 0$, find an expression for y in terms of x . [6]

Q38.

- 5 The variables x and θ satisfy the differential equation

$$2 \cos^2 \theta \frac{dx}{d\theta} = \sqrt{2x + 1},$$

and $x = 0$ when $\theta = \frac{1}{4}\pi$. Solve the differential equation and obtain an expression for x in terms of θ . [7]

Q39.

8 Let $f(x) = \frac{6 + 6x}{(2 - x)(2 + x^2)}$.

(i) Express $f(x)$ in the form $\frac{A}{2 - x} + \frac{Bx + C}{2 + x^2}$. [4]

(ii) Show that $\int_{-1}^1 f(x) dx = 3 \ln 3$. [5]

Q40.

- 2 (i) Use the trapezium rule with 3 intervals to estimate the value of

$$\int_{\frac{1}{6}\pi}^{\frac{2}{3}\pi} \operatorname{cosec} x \, dx,$$

giving your answer correct to 2 decimal places. [3]

- (ii) Using a sketch of the graph of $y = \operatorname{cosec} x$, explain whether the trapezium rule gives an overestimate or an underestimate of the true value of the integral in part (i). [2]

Q41.

- 7 In a certain country the government charges tax on each litre of petrol sold to motorists. The revenue per year is R million dollars when the rate of tax is x dollars per litre. The variation of R with x is modelled by the differential equation

$$\frac{dR}{dx} = R \left(\frac{1}{x} - 0.57 \right),$$

where R and x are taken to be continuous variables. When $x = 0.5$, $R = 16.8$.

- (i) Solve the differential equation and obtain an expression for R in terms of x . [6]
(ii) This model predicts that R cannot exceed a certain amount. Find this maximum value of R . [3]

Q42.

6 It is given that $I = \int_0^{0.3} (1 + 3x^2)^{-2} dx$.

- (i) Use the trapezium rule with 3 intervals to find an approximation to I , giving the answer correct to 3 decimal places. [3]

- (ii) For small values of x , $(1 + 3x^2)^{-2} \approx 1 + ax^2 + bx^4$. Find the values of the constants a and b .

Hence, by evaluating $\int_0^{0.3} (1 + ax^2 + bx^4) dx$, find a second approximation to I , giving the answer correct to 3 decimal places. [5]

Q43.

8 The variables x and y are related by the differential equation

$$\frac{dy}{dx} = \frac{1}{5}xy^{\frac{1}{2}} \sin\left(\frac{1}{3}x\right).$$

- (i) Find the general solution, giving y in terms of x . [6]
- (ii) Given that $y = 100$ when $x = 0$, find the value of y when $x = 25$. [3]

Q44.

10 By first using the substitution $u = e^x$, show that

$$\int_0^{\ln 4} \frac{e^{2x}}{e^{2x} + 3e^x + 2} dx = \ln\left(\frac{8}{5}\right). \quad [10]$$

[Online Classes : Megalecture@gmail.com](mailto:Megalecture@gmail.com)
www.youtube.com/megalecture
www.megalecture.com

[Online Classes : Megalecture@gmail.com](mailto:Megalecture@gmail.com)
www.youtube.com/megalecture
www.megalecture.com

[Online Classes : Megalecture@gmail.com](mailto:Megalecture@gmail.com)
www.youtube.com/megalecture
www.megalecture.com

[Online Classes : Megalecture@gmail.com](mailto:Megalecture@gmail.com)
www.youtube.com/megalecture
www.megalecture.com