

Q1.

- 2 (i) *EITHER*: Expand *RHS* and obtain at least one equation for *a* M1
 Obtain $a^2 = 9$ and $2a = 6$, or equivalent A1
 State answer $a = 3$ only A1
- OR*: Attempt division by $x^2 + ax + 1$ or $x^2 - ax - 1$, and obtain an equation in *a* M1
 Obtain $a^2 = 9$ and either $a^3 - 1 \mid a + 6 = 0$ or $a^3 - 7a - 6 = 0$, or equivalent A1
 State answer $a = 3$ only A1
- [Special case: the answer $a = 3$, obtained by trial and error, or by inspection, or with no working earns B2.] [3]
- (ii) Substitute for *a* and attempt to find zeroes of one of the quadratic factors M1
 Obtain one correct answer A1
 State all four solutions $\frac{1}{2}(-3 \pm \sqrt{5})$ and $\frac{1}{2}(3 \pm \sqrt{13})$, or equivalent A1
- [3]

Q2.

- 3 (i) Substitute $x = 3$ and equate to zero M1
 Obtain answer $\alpha = -1$ A1 2
- (ii) At any stage, state that $x = 3$ is a solution B1
EITHER: Attempt division by $(x-3)$ reaching a partial quotient of $2x^2 + kx$ M1
 Obtain quadratic factor $2x^2 + 5x + 2$ A1
 Obtain solutions $x = -2$ and $x = -\frac{1}{2}$ A1
OR: Obtain solution $x = -2$ by trial and error B1
 Obtain solution $x = -\frac{1}{2}$ similarly B2 4
- [If an attempt at the quadratic factor is made by inspection, the M1 is earned if it reaches an unknown factor of $2x^2 + bx + c$ and an equation in b and/or c .]

Q3.

- 4 (i) Substitute $x = -1$ and equate to zero obtaining e.g. $(-1)^3 - (-1)^2 + a(-1) + b = 0$ B1
 Substitute $x = 2$ and equate to 12 M1
 Obtain a correct 3-term equation A1
 Solve a relevant pair of equations for a or b M1
 Obtain $a = 2$ and $b = 4$ A1 5
- (ii) Attempt division by $x + 1$ reaching a partial quotient of $x^2 + kx$, or similar stage M1
 by inspection
 Obtain quadratic factor $x^2 - 2x = 4$ A1 2
 [Ignore failure to repeat that $x + 1$ is a factor]

Q4.

4	(i) Substitute $x = 1$ or $x = -2$ and equate to zero Obtain a correct equation, e.g. $ax + b - 5 = 0$ Obtain a second correct equation, e.g. $-8a + 4b + 4 = 0$ Solve a relevant pair of equations for a or for b Obtain $a = 2$ and $b = 3$	M1 A1 A1 M1 A1	5
	(ii) Substitute for a and b and either divide by $(x-1)(x+2)$ or attempt third factor by inspection Obtain answer $2x + 1$	M1 A1	2

Q5.

4	(i) Substitute $x = 2$, equate to zero, and state a correct equation, e.g. $16 - 12 + 2a + b = 0$ Substitute $x = -2$ and equate to -20 Obtain a correct equation, e.g. $-16 - 12 - 2a + b = -20$ Solve for a or for b Obtain $a = -3$ and $b = 2$	B1 M1 A1 M1 A1	[5]
	(ii) Attempt division by $x^2 - 4$ reaching a partial quotient of $2x - 3$, or a similar stage by inspection Obtain remainder $5x - 10$	B1 B1√ + B1√	[3]

Q6.

4	Substitute $x = -1$, equate to zero and obtain a correct equation in any form Substitute $x = -2$ and equate to 5 Obtain a correct equation in any form Solve a relevant pair of equations for a or for b Obtain $a = 2$ and $b = -3$	B1 M1 A1 M1 A1	[5]
---	--	----------------------------	-----

Q7.

6	(i) Substitute $x = 2$, equate to zero and state a correct equation, e.g. $8 + 4a + 2b + 6 = 0$ Substitute $x = 1$ and equate to 4 Obtain a correct equation, e.g. $1 + a + b + 6 = 4$ Solve for a or for b Obtain $a = -4$ and $b = 1$	B1 M1 A1 M1 A1	[5]
	(ii) <i>EITHER:</i> Attempt division by $x - 2$ reaching a partial quotient of $x^2 + kx$ Obtain remainder quadratic factor $x^2 - 2x - 3$ State linear factors $(x - 3)$ and $(x + 1)$ <i>OR:</i> Obtain linear factor $(x + 1)$ by inspection Obtain factor $(x - 3)$ similarly	M1 A1 A1 B1 B2	[3]

Q8.

- 4 (i) Commence division by $x^2 + x - 1$ obtaining quotient of the form $x + k$ M1
 Obtain quotient $x + 2$ A1
 Obtain remainder $3x + 4$ A1
 Identify the quotient and remainder correctly A1√ [4]
- (ii) Substitute $x = -1$ and evaluate expression M1
 Obtain answer 0 A1 [2]

Q9.

- 7 (i) Substitute $x = 3$ and equate to 30 M1
 Substitute $x = -1$ and equate to 18 M1
 Obtain a correct equation in any form A1
 Solve a relevant pair of equations for a or for b M1
 Obtain $a = 1$ and $b = -13$ A1 [5]
- (ii) Either show that $f(2) = 0$ or divide by $(x - 2)$, obtaining a remainder of zero B1
 Obtain quadratic factor $2x^2 + 5x - 3$ B1
 Obtain linear factor $2x - 1$ B1
 Obtain linear factor $x + 3$ B1
 [Condone omission of repetition that $x - 2$ is a factor.]
 [If linear factors $2x - 1, x + 3$ obtained by remainder theorem or inspection, award B2 + B1.] [4]

Q10.

- 4 (i) Substitute -2 and equate to zero or divide by $x + 2$ and equate remainder to zero M1
 Obtain $a = 8$ A1 [2]
- (ii) Attempt to find quotient by division or inspection or use of identity M1
 Obtain at least $3x^2 + 2x$ A1
 Obtain $3x^2 + 2x + 4$ with no errors seen A1 [3]

Q11.

- 7 (i) Substitute $x = -2$ and equate to zero M1
 Substitute $x = -1$ and equate to 24 M1
 Obtain $4a - 2b = 38$ and $a - b = 20$ or equivalents A1
 Attempt solution of two linear simultaneous equations (dependent on M1 M1) M1
 Obtain $a = -1$ and $b = -21$ A1 [5]
- (ii) Attempt to find quadratic factor by division, inspection or use of identity M1
 Obtain $6x^2 - 13x + 5$ A1√
 Conclude $(x + 2)(2x - 1)(3x - 5)$ A1 [3]

Q12.

- 3 (i) Substitute 2 and equate to zero or divide and equate remainder to zero
Obtain $a = 2$ M1
A1 [2]
- (ii) (a) Attempt to find quadratic factor by division, inspection or identity
Obtain $2x^2 + x - 3$ M1
Conclude $(x - 2)(2x + 3)(x - 1)$ A1
A1 [3]
- (b) Attempt substitution of -1 or attempt complete division by $x + 1$
Obtain 6 M1
A1 [2]

Q13.

- 3 (i) Attempt division, or equivalent, at least as far as quotient $2x + k$
Obtain quotient $2x - 3$ M1
Complete process to confirm remainder is 4 A1
A1 [3]
- (ii) State or imply $(4x^2 + 4x - 3)$ is a factor B1
Obtain $(2x - 3)(2x - 1)(2x + 3)$ B1
A1 [2]

Q14.

- 4 (i) Substitute $x = -\frac{3}{2}$, equate to zero M1
Substitute $x = -1$ and equate to 8 M1
Obtain a correct equation in any form A1
Solve a relevant pair of equations for a or for b M1
Obtain $a = 2$ and $b = -6$ A1 [5]

© Cambridge International Examinations 2013

Page 5	Mark Scheme	Syllabus	Paper
	GCE AS LEVEL – May/June 2013	9709	21

- (ii) Attempt either division by $2x + 3$ and reach a partial quotient of $x^2 + kx$, use of an identity or observation M1
Obtain quotient $x^2 - 4x + 3$
Obtain linear factors $x - 1$ and $x - 3$ A1
[Condone omission of repetition that $2x + 3$ is a factor.] A1
[If linear factors $x - 1, x - 3$ obtained by remainder theorem or inspection, award B2 + B1.] [3]

Q15.

- | | | | | |
|---|------|---|----------|-----|
| 3 | (i) | Substitute $x = -1$ and equate to zero
Obtain answer $a = 7$ | M1
A1 | [2] |
| | (ii) | Substitute $x = -3$ and evaluate expression
Obtain answer 18 | M1
A1 | [2] |

Q16.

-
- | | | | | |
|---|--|---|----------------------------|---|
| 2 | | State or obtain $-2 + a + b = 0$, or equivalent
Substitute $x = -2$ and equate to -5
Obtain 3-term equation, or equivalent
Solve a relevant pair of equations, obtaining a or b
Obtain both answers $a = 3$ and $b = -1$ | B1
M1
A1
M1
A1 | 5 |
|---|--|---|----------------------------|---|
-

Q17.

- | | | | | |
|---|----------------|---|----------------|-----|
| 3 | (i) | <i>EITHER:</i> Substitute -1 for x and equate to zero
Obtain answer $a = 6$ | M1
A1 | |
| | <i>OR:</i> | Carry out complete division and equate remainder to zero
Obtain answer $a = 6$ | M1
A1 | [2] |
| | (ii) | Substitute 6 for a and either show $f(x) = 0$ or divide by $(x - 2)$ obtaining a remainder of zero | B1 | |
| | <i>EITHER:</i> | State or imply $(x + 1)(x - 2) = x^2 - x - 2$
Attempt to find another quadratic factor by division or inspection
State factor $(x^2 + x - 3)$ | B1
M1
A1 | |
| | <i>OR:</i> | Obtain $x^3 + 2x^2 - 2x - 3$ after division by $x + 1$, or $x^3 - x^2 - 5x + 6$ after division by $x - 2$
Attempt to find a quadratic factor by further division by relevant divisor or by inspection
State factor $(x^2 + x - 3)$ | B1
M1
A1 | [4] |

Q18.

- | | | | | |
|---|--|--|----------------------------|---|
| 4 | | State or obtain $16 - 20 + 2a + b = 0$
Substitute $x = -1$ and equate to -6
Obtain a 3-term equation in any correct form
Solve a relevant pair of equations, obtaining a or b
Obtain $a = 1$ and $b = 2$ | B1
M1
A1
M1
A1 | 5 |
|---|--|--|----------------------------|---|

Q19.

2	(i) Substitute $x = 1$ and evaluate expression Obtain answer 8	M1 A1	2
	(ii) Commence division by $x^2 + x - 1$ and obtain quotient of the form $x + k$ Obtain quotient $x + 1$ Obtain remainder $2x + 4$ Correctly identify the quotient and remainder	M1 A1 A1 A1A1	4

Q20.

3	(i) Substitute $x = \frac{1}{2}$ and equate to zero Obtain answer $a = -3$	M1 A1	2
	(ii) At any stage, state that $x = \frac{1}{2}$ is a solution <i>EITHER:</i> Attempt division by $2x - 3$ reaching a partial quotient of $2x^2 + kx$ Obtain quadratic factor $2x^2 + 3x + 1$ Obtain solutions $x = -1$ and $x = -\frac{1}{2}$	M1 A1 A1	
	<i>OR:</i> Obtain solution $x = -1$ by trial and error or inspection Obtain solution $x = -\frac{1}{2}$ similarly	B1 B2	4
	[If an attempt at the quadratic factor is made by inspection, the M1 is earned if it reaches an unknown factor of $2x^2 + bx + c$ and an equation in b and/or c .]		

Q21.

5	(i) Substitute $x = -2$ and equate to zero Obtain answer $a = 3$	M1 A1	[2]
	(ii) At any stage state that $x = -2$ is a solution <i>EITHER:</i> Attempt division by $x + 2$ and reach a partial quotient of $3x^2 + kx$ Obtain quadratic factor $3x^2 + 2x - 1$ Obtain solutions $x = -1$ and $x = \frac{1}{3}$	B1 M1 A1 A1	
	<i>OR:</i> Obtain solution $x = -1$ by trial or inspection Obtain solution $x = \frac{1}{3}$ similarly	B1 B2	[4]

Q22.

2	(i) Substitute $x = -2$ and equate result to zero, or divide by $x + 2$ and equate constant remainder to zero Obtain answer $a = -13$	M1 A1	[2]
	(ii) Obtain quadratic factor $2x^2 - 5x - 3$ Obtain linear factor $2x + 1$ Obtain linear factor $x - 3$ [Condone omission of repetition that $x + 2$ is a factor.] [If linear factors $2x + 1, x - 3$ obtained by remainder theorem or inspection, award B2 + B1.]	B1 B1 B1	[3]

Q23.

- 3 (i) Substitute $x = -\frac{1}{2}$ and equate to zero M1
 Obtain $a = -11$ A1 [2]
- (ii) EITHER: Attempt division by $2x + 1$ reaching a partial quotient $2x^2 - 5x$ M1
 Obtain quadratic factor $2x^2 - 5x - 3$ A1
 Obtain complete factorisation $(2x + 1)^2(x - 3)$ A1 + A1
 OR: Obtain factor $(x - 3)$ by inspection or factor theorem B2
 Attempt division by $(x - 3)$ reaching a partial quotient $4x^2 + 4x$ M1
 Obtain complete factorisation $(2x + 1)^2(x - 3)$ A1 [4]

Q24.

- 5 (i) Substitute $x = -1$ or $x = 2$ and equate to zero M1
 Obtain a correct equation, e.g. $-a + b + 5 + 2 = 0$ A1
 Obtain a second correct equation, e.g. $8a + 4b - 10 + 2 = 0$ A1
 Solve for a or b M1
 Obtain $a = 3$ and $b = -4$ A1 [5]
- (ii) Substitute for a and b and attempt division by $(x + 1)(x - 2)$ or attempt third factor by inspection M1
 Obtain answer $3x - 1$ A1 [2]

Q25.

- 7 (i) Substitute $x = 1$, equate to zero and obtain a correct equation in any form B1
 Substitute $x = 2$ and equate to 10 M1
 Obtain a correct equation in any form A1
 Solve a relevant pair of equations for a or for b M1
 Obtain $a = -17$ and $b = 12$ A1 [5]
- (ii) At any stage, state that $x = 1$ is a solution B1
 EITHER: Attempt division by $x - 1$ and reach a partial quotient of $3x^2 + 5x$ M1
 Obtain quotient $3x^2 + 5x - 12$ A1
 Obtain solutions $x = -3$ and $x = \frac{4}{3}$ A1
- OR: Obtain solution $x = -3$ by trial and error or inspection B1
 Obtain solution $x = \frac{4}{3}$ B2

[If an attempt at the quadratic factor is made by inspection, the M1 is earned if it reaches an unknown factor of $3x^2 + 5x + \lambda$ and an equation in λ] [4]

Q26.

- 3 (i) Substitute $x = -1$ **OR** $x = 2$ correctly M1
 Equate remainders to obtain correct equation $5 - a = 26 + 2a$ or equivalent A1
 Obtain $a = -7$ A1 [3]
- (ii) Attempt division by $x - 1$ and reach a partial quotient of $x^2 + kx$ M1
 Obtain quotient $x^2 + 5x - 2$ A1
EITHER Show remainder is zero **OR** substitute $x = 1$ to obtain zero B1 [3]

Q27.

- 5 (i) Substitute $x = \frac{1}{2}$ and equate to 10 M1
 Obtain answer $a = -16$ A1
Either show that $f(3) = 0$ or divide by $(x - 3)$ obtaining a remainder of zero B1 [3]
- (ii) At any stage state that $x = 3$ is a solution B1
 Attempt division by $(x - 3)$ reaching a partial quotient of $4x^2 + kx$ M1
 Obtain quadratic factor $4x^2 - 4x - 3$ A1
 Obtain solutions $x = \frac{3}{2}$ and $x = -\frac{1}{2}$ A1
 S.C. M1A1√ if value of 'a' incorrect [4]

Q28.

- 7 (i) Substitute $x = -2$, equate to zero and obtain a correct equation in any form B1
 Substitute $x = -1$ and equate to 12 M1
 Obtain a correct equation in any form A1
 Solve a relevant pair of equations for a or b M1
 Obtain $a = 2$ and $b = 6$ A1 [5]
- (ii) Attempt division by $x + 2$ and reach a partial quotient of $2x^2 - 7x$ M1
 Obtain quotient $2x^2 - 7x + 3$ A1
 Obtain linear factors $2x - 1$ and $x - 3$ A1
 [Condone omission of repetition that $x + 2$ is a factor.]
 [If linear factors $2x - 1$, $x - 3$ obtained by remainder theorem or inspection, award B2 + B1.]
 S.C. M1A1√ if a , b not both correct [3]

Q29.

- 6 (i) Substitute $x = 1$ or $x = -2$ and equate to zero M1
 Obtain a correct equation in any form with powers of x values calculated A1
 Obtain a second correct equation in any form A1
 Solve a relevant pair of equations for a or for b M1
 Obtain $a = 3$ and $b = -5$ A1 [5]
- (ii) Attempt division by $x^2 + x - 2$, or equivalent, and reach a partial quotient of $x^2 + kx$ M1
 Obtain partial quotient $x^2 + 2x$ A1
 Obtain $x^2 + 2x - 1$ with no errors seen A1
 S.C. M1A1√ if 'a' and/or 'b' incorrect [3]

Q30.

- 7 (i) Substitute $x = -1$, equate to zero and obtain a correct equation in any form B1
 Substitute $x = 3$ and equate to 12 M1
 Obtain a correct equation in any form A1
 Solve a relevant pair of equations for a or for b M1
 Obtain $a = -4$ and $b = 6$ A1 [5]
- (ii) Attempt division by $x^2 - 2$ and reach a partial quotient of $2x - k$ M1
 Obtain quotient $2x - 4$ A1
 Obtain remainder -2 A1 [3]

Q31.

- 3 (i) Attempt division by $x^2 - 3x + 2$ or equivalent, and reach a partial quotient of $x^2 + kx$ M1
 Obtain partial quotient $x^2 - x$ A1
 Obtain $x^2 - x - 2$ with no errors seen A1 [3]
- (ii) Correct solution method for either quadratic e.g. factorisation M1
 One correct solution from solving quadratic or inspection B1
 All solutions $x = 2, x = 1$ and $x = -1$ given and no others A1 [3]

Q32.

- 4 (i) Substitute $x = 3$ and equate to 14 ($9a + 3b + 35 = 14$) M1
 Substitute $x = -2$ and equate to 24 ($4a - 2b = 24$) M1
 Obtain a correct equation in any form A1
 Solve a relevant pair of equations for a or for b M1
 Obtain $a = 1$ and $b = -10$ A1 [5]

© Cambridge International Examinations 2013

Page 5	Mark Scheme	Syllabus	Paper
	GCE AS LEVEL – October/November 2013	9709	21

- (ii) Attempt division by $x^2 + 2x - 8$ and reach a partial quotient of $x - k$ M1
 Obtain quotient $x - 1$ with no errors seen (can be done by observation) A1
 Correct solution method for quadratic e.g. factorisation M1
 All solutions $x = 1, x = 2$ and $x = -4$ given and no others CWO A1 [4]

Q33.

- 4
- (i) Substitute $x = 3$ or $x = -2$ and equate to zero M1
 Obtain a correct equation in any form A1
 Obtain a second correct equation in any form A1
 Solve a relevant pair of equations for a or for b M1
 Obtain $a = 4$ and $b = -3$ A1 [5]
- (ii) Attempt division by $x + 2$ (or $x - 3$) and obtain partial quotient of $ax^2 + kx$ M1
 Obtain linear factors $4x + 1$, $x + 2$ and $x - 3$ A1
 [If linear factor $4x + 1$ obtained by remainder theorem or inspection, award B2]
 [If linear factor $4x + 1$ obtained by division by $x^2 - x - 6$, award M1 A1] [2]
- Alternative Method:**
- Attempt to form identity $(x^2 - x - 6)(rx + s) \equiv ax^3 + bx^2 - 25x - 6$ M1
 Attempt to equate like terms M1
 Leads to $s = 1$ B1, $r = 4$ A1, $b = -3$ A1, $a = 4$ A1
 Obtain linear factors $4x + 1$, $x + 2$ and $x - 3$ A1

Q34.

- 3
- (i) Divide at least as far as x term in quotient, use synthetic division correctly or make use of an identity M1
 Obtain at least $6x^2 - x$ A1
 Obtain quotient $6x^2 - x - 2$ and confirm remainder is 7 (AG) A1 [3]
- (ii) State equation in form $(x^2 - 4)(6x^2 + kx - 2) = 0$, any constant k (may be implied) M1
 Obtain two of the roots -2 , 2 , $-\frac{1}{2}$, $\frac{2}{3}$ A1
 Obtain remaining two roots and no others A1 [3]

Q35.

- 6
- (i) Substitute -2 and equate to zero, or divide and equate remainder to zero M1
 Obtain $a = 12$ A1 [2]
- (ii) Carry out division, or equivalent, at least as far as x^2 and x terms in quotient M1
 Obtain $x^2 - 2x + 6$ A1
 Calculate discriminant of a 3 term quadratic quotient (or equivalent) DM1
 Obtain -20 (or equivalent) A1
 Conclude by referring to, or implying, root -2 and no root from quadratic factor A1 [5]

Q36.

- 5 (i) State $-40 + 4a + b = 0$ or equivalent B1
 State $-135 + 9a + b = 0$ or equivalent B1
 Solve a pair of linear simultaneous equations M1
 Obtain $a = 19$ and $b = -36$ A1 [4]
- (ii) Identify $5x - 6$ as a factor B1
 State $(x + 2)(x + 3)(5x - 6)$ B1
 State or imply $5^y = \frac{6}{5}$, following a positive value from factorisation B1√
 Apply logarithms and use power law M1
 Obtain 0.113 only A1 [5]

P3 (variant1 and 3)

Q1.

- 4 (i) Verify that $-96 + 100 + 8 - 12 = 0$ B1
 Attempt to find quadratic factor by division by $(x + 2)$, reaching a partial quotient
 $12x^2 + kx$, inspection or use of an identity M1
 Obtain $12x^2 + x - 6$ A1
 State $(x + 2)(4x + 3)(3x - 2)$ A1 [4]
 [The M1 can be earned if inspection has unknown factor $Ax^2 + Bx - 6$ and an equation in A and/or B or equation $12x^2 + Bx + C$ and an equation in B and/or C .]
- (ii) State $3^y = \frac{2}{3}$ and no other value B1
 Use correct method for finding y from equation of form $3^y = k$, where $k > 0$ M1
 Obtain -0.369 and no other value A1 [3]

Q2.

- 5 (i) Substitute $x = \frac{1}{2}$ and equate to zero, or divide, and obtain a correct equation, e.g.
 $\frac{1}{8}a + \frac{1}{4}b + \frac{5}{2} - 2 = 0$ B1
 Substitute $x = 2$ and equate result to 12, or divide and equate constant remainder to 12 M1
 Obtain a correct equation, e.g. $8a + 4b + 10 - 2 = 12$ A1
 Solve for a or for b M1
 Obtain $a = 2$ and $b = -3$ A1 [5]
- (ii) Attempt division by $2x - 1$ reaching a partial quotient $\frac{1}{2}ax^2 + kx$ M1
 Obtain quadratic factor $x^2 - x + 2$ A1 [2]
 [The M1 is earned if inspection has an unknown factor $Ax^2 + Bx + 2$ and an equation in A and/or B , or an unknown factor of $\frac{1}{2}ax^2 + Bx + C$ and an equation in B and/or C .]

Q3.

- 3 (i) Substitute $x = 2$ and equate to zero, or divide by $x - 2$ and equate constant remainder to zero, or equivalent M1
 Obtain $a = 4$ A1 [2]
- (ii) (a) Find further (quadratic or linear) factor by division, inspection or factor theorem or equivalent M1
 Obtain $x^2 + 2x - 8$ or $x + 4$ A1
 State $(x - 2)^2(x + 4)$ or equivalent A1 [3]
- (b) State any two of the four (or six) roots B1√^h
 State all roots $(\pm\sqrt{2}, \pm 2i)$, provided two are purely imaginary B1√^h [2]

Q4.

- 1 Carry out division or equivalent at least as far as two terms of quotient M1
 Obtain quotient $2x - 4$ A1
 Obtain remainder 8 A1 [3]

Q5.

- 5 (i) Substitute $x = -\frac{1}{2}$, or divide by $(2x + 1)$, and obtain a correct equation, e.g. $a - 2b + 8 = 0$ B1
 Substitute $x = \frac{1}{2}$ and equate to 1, or divide by $(2x - 1)$ and equate constant remainder to 1 M1
 Obtain a correct equation, e.g. $a + 2b + 12 = 0$ A1
 Solve for a or for b M1
 Obtain $a = -10$ and $b = -1$ A1 [5]
- (ii) Divide by $2x^2 - 1$ and reach a quotient of the form $4x + k$ M1
 Obtain quotient $4x - 5$ A1
 Obtain remainder $3x - 2$ A1 [3]

Q6.

- 10 (i) Attempt to solve for m the equation $p(-2) = 0$ or equivalent M1
 Obtain $m = 6$ A1 [2]
- Alternative:*
 Attempt $p(z) \div (z + 2)$, equate a constant remainder to zero and solve for m . M1
 Obtain $m = 6$ A1
- (ii) (a) State $z = -2$ B1
 Attempt to find quadratic factor by inspection, division, identity, ... M1
 Obtain $z^2 + 4z + 16$ A1
 Use correct method to solve a 3-term quadratic equation M1
 Obtain $-2 \pm 2\sqrt{3}i$ or equivalent A1 [5]
- (b) State or imply that square roots of answers from part (ii)(a) needed M1
 Obtain $\pm i\sqrt{2}$ A1
 Attempt to find square root of a further root in the form $x + iy$ or in polar form M1
 Obtain $a^2 - b^2 = -2$ and $ab = (\pm)\sqrt{3}$ following their answer to part (ii)(a) A1√
 Solve for a and b M1
 Obtain $\pm(1 + i\sqrt{3})$ and $\pm(1 - i\sqrt{3})$ A1 [6]

Q7.

- 3 (i) EITHER: Attempt division by $x^2 - x + 1$ reaching a partial quotient of $x^2 + kx$ M1
 Obtain quotient $x^2 + 4x + 3$ A1
 Equate remainder of form lx to zero and solve for a , or equivalent M1
 Obtain answer $a = 1$ A1
- OR: Substitute a complex zero of $x^2 - x + 1$ in $p(x)$ and equate to zero M1
 Obtain a correct equation in a in any unsimplified form A1
 Expand terms, use $i^2 = -1$ and solve for a M1
 Obtain answer $a = 1$ A1 [4]
- [SR: The first M1 is earned if inspection reaches an unknown factor $x^2 + Bx + C$ and an equation in B and/or C , or an unknown factor $Ax^2 + Bx + 3$ and an equation in A and/or B . The second M1 is only earned if use of the equation $a = B - C$ is seen or implied.]
- (ii) State answer, e.g. $x = -3$ B1
 State answer, e.g. $x = -1$ and no others B1 [2]

Q8.

7	<p>(i) Substitute $x = \frac{1}{2}$ and equate to zero</p> <p>or divide by $(2x - 1)$, reach $\frac{a}{2}x^2 + kx + \dots$ and equate remainder to zero</p> <p>or by inspection reach $\frac{a}{2}x^2 + bx + c$ and an equation in b/c</p> <p>or by inspection reach $Ax^2 + Bx + a$ and an equation in A/B</p> <p>Obtain $a = 2$</p> <p>Attempt to find quadratic factor by division or inspection or equivalent</p> <p>Obtain $(2x - 1)(x^2 + 2)$</p> <p>(ii) State or imply form $\frac{A}{2x - 1} + \frac{Bx + C}{x^2 + 2}$, following factors from part (i)</p> <p>Use relevant method to find a constant</p> <p>Obtain $A = -4$, following factors from part (i)</p> <p>Obtain $B = 2$</p> <p>Obtain $C = 5$</p>	<p>M1</p> <p>A1</p> <p>M1</p> <p>A1 cwo</p> <p>B1√</p> <p>M1</p> <p>A1√</p> <p>A1</p> <p>A1</p>	<p>[4]</p>
---	---	---	------------

Q9.

3	<p>(i) Substitute -2 and equate to zero or divide by $x + 2$ and equate remainder to zero or use -2 in synthetic division</p> <p>Obtain $a = -1$</p> <p>(ii) Attempt to find quadratic factor by division reaching $x^2 + kx$, or inspection as far as $(x + 2)(x^2 + Bx + c)$ and equations for one or both of B and C, or $(x + 2)(Ax^2 + Bx + 7)$ and equations for one or both of A and B.</p> <p>Obtain $x^2 - 3x + 7$</p> <p>Use discriminant to obtain -19, or equivalent, and confirm one root</p>	<p>M1</p> <p>A1</p> <p>M1</p> <p>A1</p> <p>A1 cwo</p>	<p>[2]</p> <p>[3]</p>
---	--	---	-----------------------

Q10.

3	<p>Substitute $x = -\frac{1}{3}$, equate result to zero or divide by $3x + 1$ and equate the remainder to zero</p> <p>and obtain a correct equation, e.g. $-\frac{1}{27}a + \frac{1}{9}b - \frac{1}{3} + 3 = 0$</p> <p>Substitute $x = 2$ and equate result to 21 or divide by $x - 2$ and equate constant remainder to 21</p> <p>Obtain a correct equation, e.g. $8a + 4b + 5 = 21$</p> <p>Solve for a or for b</p> <p>Obtain $a = 12$ and $b = -20$</p>	<p>B1</p> <p>M1</p> <p>A1</p> <p>M1</p> <p>A1</p>	<p>[5]</p>
---	---	---	------------

Q11.

- | | | | | |
|---|------|---------------|---|---------------------------------|
| 3 | (i) | <u>Either</u> | Equate $p(-1)$ or $p(-2)$ to zero or divide by $(x+1)$ or $(x+2)$ and equate constant remainder to zero.
Obtain two equations $a - b = 6$ and $4a - 2b = 34$ or equivalents
Solve pair of equations for a or b
Obtain $a = 11$ and $b = 5$ | M*1

A1
DM*1
A1 |
| | | <u>Or</u> | State or imply third factor is $4x - 1$
Carry out complete expansion of $(x+1)(x+2)(4x-1)$ or $(x+1)(x+2)(Cx+D)$
Obtain $a = 11$
Obtain $b = 5$ | B1
M1

A1
A1 [4] |
| | (ii) | | Use division or equivalent and obtaining linear remainder
Obtain quotient $4x + a$, following their value of a
Indicate remainder $x - 13$ | M1
A1 ^h
A1 [3] |

[Online Classes : Megalecture@gmail.com](mailto:Megalecture@gmail.com)
www.youtube.com/megalecture
www.megalecture.com

[Online Classes : Megalecture@gmail.com](mailto:Megalecture@gmail.com)
www.youtube.com/megalecture
www.megalecture.com