

1. (a)  $\text{Mg} + \text{H}_2\text{O} \rightarrow \text{MgO} + \text{H}_2$   
*ignore state symbols* 1  
 White solid / powder / ash / smoke  
*ignore precipitate*  
*ignore fumes* 1  
 (Bright) white light / flame  
*allow glow*  
*penalise effervescence under list principle* 1
- (b)  $2\text{Na} + \frac{1}{2}\text{O}_2 \rightarrow \text{Na}_2\text{O}$  /  $4\text{Na} + \text{O}_2 \rightarrow 2\text{Na}_2\text{O}$   
*Allow multiples, ignore state symbols*  
*Allow  $2\text{Na} + \text{O}_2 \rightarrow \text{Na}_2\text{O}_2$*  1  
 white / yellow solid / ash / smoke  
*ignore precipitate*  
*ignore fumes* 1  
 orange / yellow flame 1
2. (a)  $\text{Na}_2\text{O}$  is an ionic lattice / giant ionic / ionic crystal  
*CE=0 if molecules, atoms, metallic mentioned*  
*Mention of electronegativity max 1 out of 2* 1  
 With strong forces of attraction between ions  
*Allow strong ionic bonds / lots of energy to separate ions* 1
- (b)  $\text{SO}_3$  is a larger molecule than  $\text{SO}_2$   
*Allow greater  $M_r$  / surface area* 1  
 So van der Waals' forces between molecules are stronger  
*Any mention of ions, CE=0* 1
- (c) Ionic  
*Do not allow ionic with covalent character* 1  
 Contains  $\text{O}_2^{2-}$  ions / oxide ions  
*Equations of the form  $\text{O}_2 + \text{H}^+ \rightarrow \text{OH}^-$  /  $\text{O}_2 + 2\text{H}^+ \rightarrow \text{H}_2\text{O}$  /  $\text{O}_2 + \text{H}_2\text{O} \rightarrow 2\text{OH}^-$  score M2 and M3* 1  
 These /  $\text{O}_2$  ions (accept protons to) form OH<sup>-</sup> / hydroxide / water  
 (must score M2 to gain M3) 1
- (d) (i)  $\text{SO}_2 + \text{H}_2\text{O} \rightarrow \text{H}^+ + \text{HSO}_3^-$   
*Allow  $2\text{H}^+ + \text{SO}_3^{2-}$  but no ions, no mark*

[6]



*Only score (d)(ii) if (d)(i) correct*

- |    |      |  |  |  |      |
|----|------|--|--|--|------|
|    |      |  |  |  |      |
|    | (ii) | Reaction is an equilibrium / reversible reaction displaced mainly to the left / partially ionised / dissociated<br><i>Allow reaction does not go to completion</i> |  |  | 1    |
|    | (e)  | SiO <sub>2</sub> reacts with bases / NaOH / CaO / CaCO <sub>3</sub><br><i>Ignore incorrect formulae for silicate</i>   |  |  | 1    |
|    |      |  |  |  | 1    |
|    |      |  |  |  | [10] |
| 3. | (a)  | (i)  | Oxide 1            B   |  |      |
|    |      |  | Oxide 2 E  |  | 1    |
|    |      |  | Explanation        Low melting point or weak van der Waals' forces between molecules   |  | 1    |
|    |      | (ii)   | Chemical test Add water or flame test  |  | 1    |
|    |      |  | Test pH or flame colour  |  | 1    |
|    |      |  | Observation    pH = 13/14 or colour yellow   |  | 1    |
|    | (b)  | (i)  | Equation        CaCO <sub>3</sub> CaO + CO <sub>2</sub>  |  | 1    |
|    |      | (ii)   | Product        CaSO <sub>3</sub>   |  | 1    |
|    |      | (iii)  | Disposal of large quantities of CaSO <sub>3</sub> (allow CaSO <sub>4</sub> )   |  | 1    |
|    |      |  | Produces CO <sub>2</sub> or uses up CaCO <sub>3</sub>  |  | 1    |
|    |      |  |  |  | [10] |
| 4. | (a)  | (i)  | <i>can form a solution with pH less than 3: P<sub>4</sub>O<sub>10</sub> or SO<sub>3</sub> (1)</i>  |  |      |
|    |      | (ii)   | <i>can form a solution with with a pH greater than 12: Na<sub>2</sub>O (1)</i><br><i>penalise any wrong answer to zero</i>   |  | 2    |
|    | (b)  | (i)  | MgO + 2HNO <sub>3</sub> → Mg(NO <sub>3</sub> ) <sub>2</sub> + H <sub>2</sub> O or an ionic equation (1)<br>i.e. MgO + 2H <sup>+</sup> → Mg <sup>2+</sup> + H <sub>2</sub> O<br><i>not O<sup>2-</sup> + 2H<sup>+</sup> → H<sub>2</sub>O</i>   |  | 1    |
|    |      | (ii)   | 2NaOH + SiO <sub>2</sub> → Na <sub>2</sub> SiO <sub>3</sub> + H <sub>2</sub> O or ionic equation (1)<br>i.e. SiO <sub>2</sub> + 2OH <sup>-</sup> → SiO <sub>3</sub> <sup>2-</sup> + H <sub>2</sub> O   |  | 1    |
|    |      | (iii)  | 3Na <sub>2</sub> O + 2H <sub>3</sub> PO <sub>4</sub> → 2Na <sub>3</sub> PO <sub>4</sub> + 3H <sub>2</sub> O etc or ionic equation (1)<br>i.e. Na <sub>2</sub> O + 2H <sup>+</sup> → 2Na <sup>+</sup> + H <sub>2</sub> O  |  | 3    |
|    | (c)  |  | P <sub>4</sub> O <sub>10</sub> is a molecular (structure) or simple covalent (1)<br>Weak <u>intermolecular forces</u> or <u>van der Waals forces</u> (between molecules) (1)<br>SiO <sub>2</sub> is a macromolecule / giant covalent / giant molecule (1)<br><i>Not giant lattice</i><br>(Strong) <u>covalent</u> bonds (between atoms) must be broken (1) |  | 4    |

[9]



5. (a) Electronegativity increases 1
- Proton number increases (increase in nuclear charge) 1
- Same number of electron shells/levels 1  
*Or same radius or Shielding of outer electrons remains the same*
- Attraction of bond pair to nucleus increases 1  
*Allow 'electrons in bond' instead of 'bond pair'* 1

[www.megalecture.com](http://www.megalecture.com)



- (b) Big difference in electronegativity leads to ionic bonding,  
smaller covalent  
*Lose a mark if formula incorrect* 1
- Sodium oxide ionic lattice 1
- Strong forces of attraction between ions 1
- $P_4O_{10}$  covalent molecular 1  
*Must have covalent and molecular (or molecules)*
- Weak (intermolecular) forces between molecules 1  
*Or weak vdW, or weak dipole–dipole between molecules*
- melting point  $Na_2O$  greater than for  $P_4O_{10}$  1  
*Or argument relating mpt to strength of forces*
- (c) Moles  $NaOH = 0.0212 \times 0.5 = 0.0106$  1  
*M1 moles of NaOH correct*
- Moles of  $H_3PO_4 = 1/3$  moles of  $NaOH (= 0.00353)$  1  
*M2 is for 1/3*
- Moles of P in 25000 l =  $0.00353 \times 10^6 = 3.53 \times 10^3$  1  
*M3 is for factor of 1,000,000*
- Moles of  $P_4O_{10} = 3.53 \times 10^3/4$  1  
*M4 is for factor of 1/4 (or 1/2 if  $P_2O_5$ )*
- Mass of  $P_4O_{10} = 3.53 \times 10^3/4 \times 284 = 0.251 \times 10^6$  g 1  
 $= 251$  kg  
*(Or if  $P_2O_5$   $3.53 \times 10^3/2 \times 142$ )*  
*M5 is for multiplying moles by M, with correct units*  
*allow conseq on incorrect M4*  
*(allow 250-252)*

[15]