



Topic 2 Exercise 3 - Ideal Gas Equation

Remember: $R = 8.31 \text{ JK}^{-1}\text{mol}^{-1}$, $0 \text{ K} = -273 \text{ }^{\circ}\text{C}$

- Calculate the volume occupied by one mole of a gas at $25 \text{ }^{\circ}\text{C}$ and 100 kPa .
- Calculate the pressure of a gas given that 0.2 moles of the gas occupy 10 dm^3 at $20 \text{ }^{\circ}\text{C}$.
- Calculate the temperature of a gas if 0.5 moles occupy 1.2 dm^3 at a pressure of 200 kPa .
- Calculate the mass of a sample of carbon dioxide which occupies 20 dm^3 at $27 \text{ }^{\circ}\text{C}$ and 100 kPa .
- Calculate the relative molecular mass of a gas if a 500 cm^3 sample at $20 \text{ }^{\circ}\text{C}$ and 1 atm has a mass of 0.66 g .
- At $25 \text{ }^{\circ}\text{C}$ and 100 kPa a gas occupies a volume of 20 dm^3 . Calculate the new temperature of the gas if
 - the volume is decreased to 10 dm^3 at constant pressure.
 - the pressure is decreased to 50 kPa at constant volume.
- 10.0 g of calcium nitrate is heated at 100 kPa and a temperature of $300 \text{ }^{\circ}\text{C}$, at which temperature it fully decomposes. Calculate
 - the volume of nitrogen dioxide evolved
 - the volume of oxygen evolved
 - the total volume of gas evolved

Equation: $2\text{Ca}(\text{NO}_3)_2(\text{s}) \rightarrow 2\text{CaO}(\text{s}) + 4\text{NO}_2(\text{g}) + \text{O}_2(\text{g})$
- Calculate the volume of oxygen produced at 298 K and 100 kPa by the decomposition of 30 cm^3 of 0.1 mol dm^{-3} hydrogen peroxide.

Equation: $2\text{H}_2\text{O}_2(\text{aq}) \rightarrow 2\text{H}_2\text{O}(\text{l}) + \text{O}_2(\text{g})$
- Lead (IV) oxide dissolves in concentrated hydrochloric acid according to the following equation: $\text{PbO}_2(\text{s}) + 4\text{HCl}(\text{aq}) \rightarrow \text{PbCl}_2(\text{s}) + \text{Cl}_2(\text{g}) + 2\text{H}_2\text{O}(\text{l})$
Starting with 37.2 g of lead (IV) oxide, calculate:
 - the volume of 12 mol dm^{-3} HCl needed to completely dissolve it
 - the mass of PbCl_2 produced
 - the volume of chlorine produced at 298 K and 100 kPa .
- What mass of magnesium, and what volume of 2.0 mol dm^{-3} hydrochloric acid, will be required to produce 100 cm^3 of hydrogen gas at 298 K and 100 kPa ?

Equation: $\text{Mg}(\text{s}) + 2\text{HCl}(\text{aq}) \rightarrow \text{MgCl}_2(\text{aq}) + \text{H}_2(\text{g})$
- 0.52 g of sodium was added to 100 cm^3 of water. Calculate:
 - The volume of hydrogen evolved at 298 K and 100 kPa
 - The concentration of the sodium hydroxide solution produced, assuming the volume of water does not change.

Equation: $2\text{Na}(\text{s}) + 2\text{H}_2\text{O}(\text{l}) \rightarrow 2\text{NaOH}(\text{aq}) + \text{H}_2(\text{g})$