


1. **QoL** Bonding Both covalent
(linked statement) 1
- Structure Iodine = molecular /I₂ (stated or in diagram)
[treat incorrect diagram as contradiction] 1
- Diamond = giant molecular/macromolecular/giant
covalent / giant atomic (stated only)
Reference to van der Waals' /dipole-dipole = contradiction 1
- QoL** Iodine Weak van der Waals' forces / induced dipole-induced
dipole 1
- Diamond Covalent bonds would need to be broken 1
- Many / strong covalent bonds **OR** much energy needed
Tied to M5 or near miss
[If ionic/metallic structure suggested then CE for that
substance]
[If hydrogen bonding suggested, for I₂ lose M2 & M4; for
diamond lose M3, M5 & M6] 1
2. (a) Ability/power of an atom/element/nucleus to withdraw electron
density or electron cloud or a pair of electrons (towards itself);
Not withdraw an electron
If ref to ionic, metallic, IMF etc then CE = 0 1
- From a covalent bond or from a shared pair of electrons;
Not distort
Not remove electrons 1
- (b) Van der Waals/ vdw/London/ temporary (induced) dipole/
dispersion forces; 1
- Hydrogen bonds/H bonds;
Not just hydrogen 1

[6]

- (c) (Large) electronegativity difference between N + H/ difference of 0.9/ N very electronegative;
Insufficient to say $N = 3.1$ and $H = 2.1$ 1
- Forms N $-$ / H $+$ or dipole explained in words;
Not N becomes (fully) negative or vice versa 1
- Lone pair on N attracts/forms weak bonds with H ($+$);
 QWC
Can score M2 and 3 from a diagram 1
- (d) Co-ordinate/dative;
If not correct then CE = 0. If covalent/blank mark on. 1
- Both electrons/ lone pair (on P/PH₃)
Not lone pair on hydrogen 1
- Shares/donated from P(H₃)/ to H($+$); 1
- (e) 3 bonds and 1 lp attached to As;
Must label H and As atoms
Accept distorted tetrahedral not bent tetrahedral 1
- Pyramidal/tetrahedral/ trigonal pyramidal;
Not bipyramidal/triangular 1
- (f) (Only) weak Van der Waals forces between molecules /AsH₃ has weaker IMF /ammonia has hydrogen bonding/ more energy needed to break IMF's in ammonia/ Van der Waals weaker than H bonds;
Accept has no H bonds.
Ignore dp-dp in AsH₃ provided ammonia has stronger IMF.
If between atoms mentioned CE=0
Break bonds CE = 0 1
- (g) $4\text{AsCl}_3 + 3\text{NaBH}_4 \rightarrow 4\text{AsH}_3 + 3\text{NaCl} + 3\text{BCl}_3$;
Accept multiples 1

[14]

3. (a) (i)



M1 Si: cross 1200 1

M2 Cl: cross below S 1

M3 Ar: cross below Cl
[allow, even if M2 wrong]
[If Cl cross missing and Ar below S, allow M3] 1

(ii) Si is macromolecular/giant molecular/giant covalent/ giant atomic 1

Covalent bonds need to be broken/accept 'overcome'
[Not loosened/weakened] 1

Covalent bonds are strong / many covalent bonds involved/
 requires much energy/hard to break
[Tied to 'break' or near miss in M2] [Not 'structure' is broken]
[Must mention 'covalent' somewhere in part (a)(ii) to earn M2/M3]
[If van der Waals'/IMF mentioned M2/M3 = CE = 0.]
[If ions mentioned M1/M2/M3 = CE = 0] 1

(iii) Intermolecular force = van der Waals'/induced dipole-dipole/dispersion forces 1

QoL Sulphur has greater M_r / size / surface area/more electrons/more atoms **so** stronger intermolecular forces (comparison)
[Mark separately] [Not 'more shells'] 1

(b) Trend: Decreases
 [If trend wrong = CE = 0] 1

Increase in size of ion/atom / more shells / decrease in charge density / decrease in charge size ratio 1

Weaker attraction for delocalised/free/sea of electrons / weaker metallic bonding
 [Ignore shielding]
 [van der Waals' etc. = CE = 0 for M2 and M3] 1

[11]

4. (a)



[Diagrams must be complete and accurate]

2

(b) (i) Attraction /electrostatic forces/bonds/attractions between (positive) ions/lattice and delocalised/free electrons/sea of electrons.
 [Not metallic bonding]
 [Not just 'forces'] 1

(ii) Electrostatic attractions/forces between ions or attractions between (oppositely charged) ions/ Na⁺ & Cl⁻
 [Not ionic bonding] 1

(iii) (Here) the ionic bonding in NaCl is stronger/requires more energy to break than the metallic bonding in Na

QoL Accept 'bonding/forces of attraction in NaCl is stronger than in Na'
 [If IMF/molecules/van der Waals'/dipole-dipole mentioned in parts(i) or (ii), then CE = 0 for parts (i) and/or(ii) and CE = 0 for part(iii)] 1

- (c) Comparison:
Sodium conducts **and** sodium chloride does NOT conduct
Allow 'only Na conducts'
Accept 'Na conducts, NaCl only conducts when molten'
[Do not accept sodium conducts better than sodium chloride etc.]

1

Explanation:
(Delocalised) electrons flow through the metal

1

Allow e^- move/carry current/are charge carriers/transfer charge.

[Not 'electrons carry electricity']

[Not 'NaCl has no free charged particles']

Ions can't move in solid salt

1

- (d) Layers can slide over each other – idea that ions/atoms particles move
[Not molecules]
[Not layers separate]

1

[12]

5. (a) Oxygen more/very/highly electronegative (than hydrogen)
OR oxygen has stronger attraction for bonding electrons / bonding electrons drawn towards oxygen;

1

causes higher e^- density round oxygen atom / causes H^+
 O^- ;

1

- (b) van der Waals' forces between oxygen molecules;

1

Hydrogen bonding between methanol molecules;

1

H-B stronger than van der Waals' OR stronger IMF in methanol;

(if dipole-dipole forces in O_2 or methanol, allow comparison, hence max 2)

(if ionic/covalent etc. max 1)

(mention of bond break = CE = 0)

1

[5]



- | | | |
|-----|---|-----|
| 6. | B | [1] |
| 7. | C | [1] |
| 8. | C | [1] |
| 9. | A | [1] |
| 10. | B | [1] |