

## AS LEVEL CHEMISTRY

## TOPIC 8 – REACTIONS OF ORGANIC COMPOUNDS TEST

Answer all questions

Max 50 marks

Name ......./50 ......% Grade .......

Page 1 of 10

## MEGA LECTURE

- 1. Trifluoromethane (CHF<sub>3</sub>) can be used to make the reinigerant chilorotrifluoromethane(CCIF<sub>3</sub>).
  - (a) Chlorotrifluoromethane is formed when trifluoromethane reacts with chlorine.

(b)

CHF<sub>3</sub> + Cl<sub>2</sub> CCIF<sub>3</sub> + HCI

The reaction is a free-radical substitution reaction similar to the reaction of methane with chlorine.


(i)	Write an equation for each of the following steps in the mechanism for the reaction of $\text{CHF}_{\scriptscriptstyle 3}$ with $\text{Cl}_{\scriptscriptstyle 2}$	
	Initiation step	
	First propagation step	
	Second propagation step	
	Termination step to form hexafluoroethane	
		(4)
(ii)	Give <b>one</b> essential condition for this reaction.	
		(1)
	ome refrigeration systems, CHF₃ has replaced CCIF₃ because of concerns about ozone etion.	
(i)	Identify the species formed from CCIF <sub>3</sub> that is responsible for the catalytic decomposition of ozone in the upper atmosphere.	
		(1)
(ii)	Write an overall equation to represent the decomposition of ozone into oxygen.	

(Total 7 marks)

2. Consider the following reactions



 $H_2C = CHCH_3$   $\longrightarrow$   $CH_3CHBrCH_3$   $\longrightarrow$   $CH_3CH(NH_2)CH_3$   $\longrightarrow$   $NH_3$ 



substance X

(a) Name and outline a mechanism for Reaction 1.

Name of mechanism .....

Mechanism

(5)

(b) Name and outline a mechanism for Reaction 2.

Name of mechanism

Mechanism



(C)	Give the name of substance <b>X</b> .		
		(2)	
(d)	The haloalkane produced in Reaction 1 can be converted back into propene in an elimination reaction using ethanolic potassium hydroxide.	( )	

CH<sub>3</sub>CHBrCH<sub>3</sub>  $\longrightarrow$  H<sub>2</sub>C=CHCH<sub>3</sub>

Outline a mechanism for this conversion.

(3) (Total 15 marks)

3. Glucose can be used as a sour MEGA LECTURE as a fuel or can be converted into ethene.

		$C_6H_{12}O_6$	CH₃CH₂OH	H <sub>2</sub> C=CH <sub>2</sub>	
		glucose	ethanol	ethene	
(a) I	Nam	e the types	of reaction illustr	ated by the two reactions above.	
(	Gluc	ose to ethai	nol		
ı	Etha	nol to ether	ne		
(b) (	(i)	State what	t must be added t	to an aqueous solution of glucose so that ethanol is formed.	(2
(	(ii)	Identify a s	suitable catalyst fo	or the conversion of ethanol into ethene.	10
(c) (	(i)		class of alcohols t	to which ethanol belongs.	(2
(	(ii)	Give <b>one</b> a fraction.	advantage of usin	ng ethanol as a fuel compared with using a petroleum	
(d) I	Most abse	t of the ethe ence of air. V	ne used by indus Vrite an equation	stry is produced when ethane is heated to 900°C in the for this reaction.	(2
'a\ .		o the tune o		which accurs when others is converted into poly(others)	(1
(e) I	ivam	ie irie type c	or polymensation	which occurs when ethene is converted into poly(ethene).	
					(1 arks

acs	app	• Гс	anad Hameed +92 323 509 4443, email: megalecture@gmail.c			
4.	In ar pota	an investigation of the chemi MEGA LECTURE of ethanol and acidified tassium dichromate(VI) is he				
	(a)	Ехр	lain why a water bath is used to heat the mixture.			
			(1)			
	(b) Describe the colour change which would be observed.					
		•••••				
			(1) (Total 2 marks)			
5.	(a)	Ethanol can be manufactured by the direct hydration of ethene and by the fermentation of sugars.				
		(i)	State what is meant by the term <i>hydration</i> .			
		(ii)	Give <b>one</b> advantage and <b>one</b> disadvantage of manufacturing ethanol by fermentation rather than by hydration.			
			Do <b>not</b> include energy consumption or cost.			
			Advantage			
			Disadvantage			

(3)

- (b) Ethanol can be oxidised t MEGA LECTURE
  - (i) Draw the structure or this aluenyoe and or this carboxylic acid.

Structure of aldehyde

Structure of carboxylic acid

(ii)	Give a suitable reagent and reaction conditions for the oxidation of ethanol to form the carboxylic acid as the major product.				
	Reagent				
	Conditions				
		<i>7</i> , •			
<i>a</i> .		~~~			

(c) (i) Draw the structure of an alcohol containing four earbon atoms which is resistant to oxidation.

(ii) Draw the structure of an alcohol containing four carbon atoms which can be oxidised to a ketone.

(2) (Total 10 marks)

(5)

- 6. Which one of the following med MEGA LECTURE on sequence below CH<sub>3</sub>CH<sub>3</sub> CH<sub>3</sub>CH<sub>2</sub>CI CH<sub>3</sub>CH<sub>2</sub>CH CH<sub>3</sub>CH CH<sub>3</sub>CH
  - A electrophilic addition
  - B electrophilic substitution
  - C nucleophilic substitution
  - **D** free-radical substitution

В

C

D

(Total 1 mark)

7. In which of the following is a curly arrow used incorrectly?

$$CH_3CH_2CHCH_3 \longrightarrow CH_3CH_2CHCH_3 + :Br - OH$$

$$CH_3CH_2CCH_3 \longrightarrow CH_3CH_2CCH_3 \longrightarrow CH_3CH_2CCH_3$$

$$\downarrow \\ :NH_3 \qquad \qquad \downarrow \\ H \longrightarrow NH_2 \qquad \qquad NH_2$$

$$CH_3CH_2CHCH_3 \longrightarrow CH_3CH \longrightarrow CH_3CH = CHCH_3$$

(Total 1 mark)

- **8.** Which one of the following statements explains best why fluoroalkanes are the least reactive haloalkanes?
  - **A** Fluorine is much more electronegative than carbon.
  - **B** The F<sup>-</sup> ion is the most stable halide ion.
  - **C** The C–F bond is the most polar carbon–halogen bond.
  - **D** The C–F bond is the strongest carbon–halogen bond.

(Total 1 mark)

whatsapp: Fahad Hameed +92 323 509 4443, email: megalecture@gmail.com MEGA LECTURE Which one of the following con-9. Α propene propane В propan-I-ol propanal C propanal propanoic acid D propanone propane (Total 1 mark) Which one of the following isomers is not oxidised under mild reaction conditions? 10. Α (CH<sub>3</sub>)<sub>2</sub>CHCH(OH)COCH<sub>3</sub> В (CH<sub>3</sub>)<sub>2</sub>C(OH)CH<sub>2</sub>COCH<sub>3</sub> C (CH<sub>3</sub>)<sub>2</sub>CHCH(OH)CH<sub>2</sub>CHO D (CH<sub>3</sub>)<sub>2</sub>C(OH)CH<sub>2</sub>CH<sub>2</sub>CHO (Total 1 mark) Which one of the following alcohols forms a mixture of alkenes when dehydrated? of Α propan-1-ol В propan-2-ol C pentan-1-ol D pentan-2-ol (Total 1 mark) Which one of the following cannot be produced by oxidation of propan-I-ol? 12. Α carbon dioxide В propanone C propanal D propanoic acid (Total 1 mark)

 25.0 cm³ of ethanedioic acid red complete neutralisation.



The concentration of ethanedioic acid is

- **A** 0.0225 mol dm<sup>-3</sup>
- **B** 0.0450 mol dm<sup>-3</sup>
- **C** 0.0560 mol dm<sup>-3</sup>
- **D** 0.0900 mol dm<sup>-3</sup>

(Total 1 mark)